



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/645, 31/55	A1	(11) International Publication Number: WO 99/59598 (43) International Publication Date: 25 November 1999 (25.11.99)
(21) International Application Number: PCT/CA99/00449 (22) International Filing Date: 19 May 1999 (19.05.99) (30) Priority Data: 09/081,709 19 May 1998 (19.05.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/081,709 (CIP) Filed on 19 May 1998 (19.05.98) (71) Applicant (for all designated States except US): DALHOUSIE UNIVERSITY [CA/CA]; Office of the President, Arts and Administration Building, 6299 South Street, Halifax, Nova Scotia B3H 4H6 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): SAWYNOK, Jana [CA/CA]; 57 Foster Avenue, Fall River, Nova Scotia B2T 1E7 (CA). ESSER, Mike [CA/CA]; 117 Johnson Avenue, Dartmouth, Nova Scotia B2Y 2K8 (CA). REID, Allison [CA/CA]; 202-1326 Lower Water Street, Halifax, Nova Scotia B3J 3R3 (CA).		(74) Agents: MORROW, Joy, D. et al.; Smart & Biggar, 900-55 Metcalfe Street, P.O. Box 2999, Station D, Ottawa, Ontario K1P 5Y6 (CA). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: USE OF TRICYCLIC ANTIDEPRESSANTS FOR LOCAL ANALGESIA (57) Abstract <p>When administered locally, tricyclic, second generation and third generation antidepressants, such as amitriptyline and desipramine, have been shown to produce analgesia in a subject having a site of local discomfort. The analgesic effect of such antidepressants, when administered locally is equal to that achieved by systemic administration and lasts longer. The invention provides compositions containing tricyclic, second generation and third generation antidepressants for local administration, such as those formulated for topical application, or for injection in slow release delivery vehicles; and methods for their use for producing local analgesia.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF TRICYCLIC ANTIDEPRESSANTS FOR LOCAL ANALGESIA

FIELD OF THE INVENTION

5 The present invention relates to therapeutic compositions and methods for their use. In a particular aspect, the present invention relates to compositions for producing pain relief and methods of their use.

BACKGROUND OF THE INVENTION

10

Neuropathic pain is a form of chronic pain that can persist for months, years or decades following an injury, and results from damage to peripheral nerves, nerve roots, the spinal cord or certain brain regions. It differs from nociceptive pain in terms of duration, characteristics, underlying mechanisms and treatment (Bennett G.J. (1994a) In: *Textbook of Pain* (Ed. Wall PD, Melzack R) Churchill Livingstone, London 3rd edn, 201). Neuropathic pain can consist of spontaneous pain (e.g., burning, cutting, tingling), evoked pain (e.g., allodynia evoked by stimulation of non-nociceptive afferents, and hyperalgesia evoked by stimulation of nociceptive afferents) and paroxysmal pain (e.g., originating from a trigger point, described as

15 stabbing, lancinating, shocklike) (Bennett G.J. (1994a) In: *Textbook of Pain* (Ed. Wall P.D., Melzack R.) Churchill Livingstone, London 3rd edn, 201). Neuropathic pain can accompany nociceptive pain, and multiple treatment strategies may be required for optimal alleviation of pain (Portenoy R.K. (1991) In: *Towards a New Pharmacology of Pain* (Ed. Basbaum A.I., Besson J.M.) John Wiley & Sons Ltd, New

20 York, 393; Devor M. et al. (1991) In: *Towards a New Pharmacotherapy of Pain* (Ed. Basbaum A.I., Besson J.M.) John Wiley & Sons, New York, 417). Neuropathic syndromes are traditionally classified according to the disease or event that precipitated them (e.g., postherpetic neuralgia following shingles, causalgia following partial damage to a major nerve, central pain following a thalamic infarct) (Portenoy

25 R.K. (1991) In: *Towards a New Pharmacology of Pain* (Ed. Basbaum A.I., Besson J.M.) John Wiley & Sons Ltd, New York, 393; Merskey H., Bogaduk N. (1994) In:

30

- Classification of Chronic Pain. Descriptions of Chronic Pain Syndromes and Definitions of Pain Terms*, 2nd edn., IASP Press, Seattle, page 40). The involvement of the sympathetic nervous system in a number of these conditions has been appreciated for some time. Pain syndromes with a sympathetic component are considered as sympathetically maintained pain (reflect sympathetic dystrophy, causalgia) (Bonica J.J. (1990) In: *The Management of Pain* (Ed. Bonica J.J.) Lea & Fibiger, Philadelphia 2nd edn., 220; Blumberg H., Jänig W. (1994) In: *Textbook of Pain* (Ed. Wall P.D., Melzack R.) Churchill Livingstone, London 3rd edn., 685).
- 10 Neuropathic pain is particularly difficult to treat clinically. The use of opioids is controversial, with issues of contention being the relative refractoriness of neuropathic pain compared to nociceptive pain, the need for higher doses with an increased incidence of side effects to achieve partial effects, and concerns over the long term use of opioids in a non-malignant context (Arnér S., Meyerson B.A. (1988) *Pain* 22: 11; Kuypers H., König H., Adriaensson H., Gybels J.M. (1991) *Pain* 47: 5; Portenoy R.K., Foley K.M., Inturrisi C.E. (1990) *Pain* 43: 273; Portenoy R.K. (1994) In: *Progress in Pain Research and Management* (Ed. Fields H.L., Liebskind J.C.) IASP Press, Seattle, 247). The major classes of agents currently used to treat neuropathic pain include systemically delivered antidepressants, anticonvulsants, local anesthetics, and specialized agents such as muscle relaxants, and sympatholytic drugs
- 15 (reviewed Portenoy R.K. (1991) In: *Towards a New Pharmacology of Pain* (Ed. Basbaum A.I., Besson J.M.) John Wiley & Sons Ltd, New York, 393; Portenoy R.K. (1993) *Drug Therapy* 23: 41; Max M.B. (1994) In: *Progress in Pain Research and Management* (Ed. Fields H.L., Liebskind J.C.) IASP Press, Seattle, 229). However,
- 20 many of these treatments show limited effectiveness (complete pain relief is rarely achieved), and there is a high incidence of debilitating side effects (Portenoy R.K. (1993) *Drug Therapy* 23: 41; Bennett G.J. (1994a) In: *Textbook of Pain* (Ed. Wall P.D., Melzack R.) Churchill Livingstone, London 3rd edn, 201; Mac Farlane et al., (1997) *Pharmacol. Ther.* 75:1).
- 25

Damage to nerves can activate changes in both the peripheral and central nervous systems, and these lead to the characteristic expression of neuropathic pain. Peripherally, the following mechanisms may be involved (reviewed by Devor M et al. (1991) In: *Towards a New Pharmacotherapy of Pain* (Ed. Basbaum A.I., Besson J.M.) John Wiley & Sons, New York, 417; Devor M. (1994) In: *Textbook of Pain* (Ed. Wall P.D., Melzack R.) Churchill Livingstone, London, 3rd edn., 79; Blumeberg H., Jänig W. (1994) In: *Textbook of Pain* (Ed. Wall P.D., Melzack R.) Churchill Livingstone, London 3rd edn, 685). (1) Following injury, normally silent nociceptors become sensitized and acquire an ongoing spontaneous activity, a lowered threshold for activation, and a heightened response to suprathreshold stimulation. (2) Development of ectopic discharges can occur at the site of injury, site of regeneration and neuroma formation, or within the dorsal root ganglion. (3) Following injury, sympathetic afferents can excite sensory nerve endings, regenerating nerve sprouts, neuromas, and the dorsal root ganglion cell. Central mechanisms implicated in neuropathic pain are as follows (Coderre et al., (1993) *Pain* 52:259; Bennett (1994) In: *Textbook of Pain* (ed. by P.D. Wall et al., London, 3rd edn., 201; Woulff and Doubell (1994) *Curr. Opin. Neurobiol.* 4:525): (1) Transmission neurons within the spinal cord exhibit enhanced sensitivity to excitation by excitatory amino acids (EAAs) and substance P (SP). This condition leads to allodynia (following A β fibre activation) and hyperalgesia (following A δ and C fibre activation). (2) Disinhibition resulting from loss of inhibitory input to projection neurons and an aberrant patterning of responses to large diameter afferent activation. (3) Myelinated afferents, which normally would enervate deeper laminae, projecting into and making synaptic connections in more superficial laminae. This can be a mechanism for the generation of allodynia.

A number of models of neuropathic pain have been developed in order to provide a basis for understanding of neuropathic changes that occur following nerve

injury, and to provide model systems in which to test novel therapeutic strategies (reviewed Bennett G.J. (1994b) In: *Progress in Pain Research and Management* Vol 2 (Ed. Gebhart C.F., Hammond D.L., Jensen T.S.) IASP Press, Seattle, 495). The best characterized are a number of nerve injury models which have been developed

5 only relatively recently (chronic constriction injury due to loose ligation of the sciatic nerve, Bennett G.J., Xie Y.K. (1988) *Pain* 33: 87; and partial sciatic ligation due to tight ligation of a portion of the sciatic nerve (Seltzer et al. (1990) *Pain* 43:205) and tight ligation of two spinal nerves (Kim S.H., Chung J.M. (1992) *Pain* 50: 355)).

These model systems exhibit various manifestations of neuropathic pain (e.g., degree

10 of expression of spontaneous pain behaviors, mechanical or thermal allodynia, mechanical or thermal hyperalgesia) and degree of sympathetic nerve involvement (Neil A., et al. (1991) *Brain Res* 565: 237; Shir Y., Seltzer Z. (1991) *Pain* 45: 309; Kim S.H., Na H.S., Sheen Ki, Chung J.M. (1993) *Pain* 55: 85; Kim K.J., Yoon Y.W., Chung J.M. (1997) *Exp Brain Res* 113: 200). The spinal nerve ligation (SNL) model

15 exhibits a greater degree of evoked pain (mechanical allodynia) and of sympathetic involvement in this parameter than do the partial sciatic ligation or chronic constriction injury models, while the chronic constriction injury model exhibits a greater degree of spontaneous pain. It is however appreciated that such properties may be relative and change with time (i.e., sympathetic dependency can change with

20 time both in animal models and in clinical neuropathic pain, Staton-Hicks M, et al. (1995) *Pain* 63: 127).

Systemically administered antidepressants offer an alternate therapy in neuropathic and chronic pain states. Interactions with biogenic amines, endogenous

25 opioids, excitatory amino acid receptors, substance P and calcium and sodium channels have been considered in efforts to pinpoint the mechanism of systemically administered antidepressants (reviewed by Eschalier A. et al. (1994) *CNS Drugs* 2: 261). What is clear is that antidepressants can act at both supraspinal (Spiegel, K. et al., *Ann. Neurol.* 13 (1983) 462-465, Eschalier A, et al. (1994) *CNS Drugs* 2: 261).

and spinal (Hwang, A.S. and Wilcox, G.L., *Pain* 28 (1987) 343-355; Iwashita, T. and Shimizu, T., *Brain Research* 581 (1992) 59-66; Eisenachand J. and Gebhart G.F., *Anesthesiology* 83 (1995) 1046-1054) sites of action.

5 This analgesic action is independent of antidepressant effects as it occurs in non-depressed subjects and occurs independently of mood changes in depressed subjects (Magni G. (1991) *Drugs* 42: 730; Onghena P. and Van Houdenhove B (1992) *Pain* 49: 205; Max M.B. (1994) In: *Progress in Pain Research and Management* (Ed. Fields H.L., Liebskind J.C.) IASP Press, Seattle, 229, McQuay H.J., Tramer M., Nye
10 B.A., Carroll D., Wiffen P.J., Moore R.A. (1996) *Pain* 68: 217). Agents which block the uptake of both noradrenaline (NA) and 5-hydroxytryptamine (5-HT) such as amitriptyline, or which block NA but not 5-HT, such as desipramine, are more effective than those with selectivity for 5-HT, such as fluoxetine (Max M.B. (1994) In: *Progress in Pain Research and Management* (Ed. Fields H.L., Liebskind
15 J.C.) IASP Press, Seattle, 229). Pain relief is reported to be apparent within one week of therapy (McQuay H.J., Carroll D., Glynn C.J. (1992) *Anaesthesia* 47: 646). This time course corresponds to the time required to attain stable plasma levels ($t_{1/2}$ 17-36 hours in humans, Ziegler V.E., Biggs J.T., Aardekani A.B., Rosen S.H. (1978) *J Clin Pharmacol* 18: 462). By contrast, the antidepressant activity of these compounds
20 takes 4-6 weeks to become apparent (Potter W.Z., Rudorfer M., Manji H. (1991) *New Eng J Med* 325: 633). These differences in profile of active drugs, time course, and independent expression of effects suggest that mechanisms underlying pain relief and alleviation of depression differ.

25 In animal tests, both the systemic and spinal administration of antidepressants show intrinsic efficacy in a number of nociceptive pain tests, and augment analgesia produced by opioids (reviewed by Eschalier A, Mestre C, Dubray C, Ardid D (1994) *CNS Drugs* 2: 261). However, this profile can be variable, and inhibitory effects on the action of morphine have been observed in some cases (reviewed by Eschalier et

- al., supra). Methodological issues (e.g., test paradigm, intensity of stimulus, dose, regimen of acute versus chronic administration) are reported to account for many of these differences (Kellstein D.E., Malseed R.T., Goldstein F.J. (1984) *Pain* 60: 275; Kellstein D.E., Malseed R.T., Ossipov M.H., Goldstein F.J. (1988) *Neuropharmacology* 27: 1; Fialip J., Marty H., Makambila M.C., CiViate M.A., Eschalier A. (1989) *J Pharmacol Exp Ther* 248: 747). Systemically administered antidepressants also exhibit intrinsic actions in a number of neuropathic pain tests including nerve transaction (Seltzer Z., Tal M., Sherav Y. (1989) *Pain* 37: 245), mononeuropathy (Ardid D, Gilbaud G (1992) *Pain* 49: 279) and diabetic neuropathy models (C.Courteix et al. (1994) *Pain* 57:153-160). One study examined chronic versus acute dosing regimens (Ardid D., Gilbaud G. (1992) *Pain* 49: 279), and observed that the activity seen following chronic paradigms appeared to be accounted for by accumulating doses rather than being qualitatively different.

15

SUMMARY OF THE INVENTION

- In accordance with the present invention, it has been discovered that locally administered tricyclic, second generation, or third generation antidepressant(s) produce a local antinociceptive action, especially against inflammatory and neuropathic pain. When administered locally in animal models of inflammatory (formalin test) and neuropathic pain (spinal nerve ligation), amitriptyline, a non selective noradrenaline (NA) and 5-hydroxytryptamine (5-HT) reuptake inhibitor, and desipramine, a selective NA reuptake inhibitor, produced local antinociceptive actions.

25

- Accordingly, in one aspect of the invention, there are provided compositions containing an effective therapeutic amount of a tricyclic, second generation or third generation antidepressant in an inert carrier. The compositions are preferably formulated for local administration, for example in a saline solution, or as a cream, gel, ointment, or spray.

30

Such formulations possess the advantage that a higher and more efficacious concentration of the antidepressant can be attained in the region of the sensory nerve terminal than with systemic administration of the antidepressant. In addition, local
5 administration greatly reduces the side effects that result from systemic administration of tricyclic, second generation or third generation antidepressant(s).

In another aspect of the invention, there are provided methods for producing local analgesia in a subject having a site of local discomfort, such as is caused by
10 inflammatory or neuropathic conditions. In the invention method, local analgesia is obtained by locally administering an effective amount of a tricyclic, second generation or third generation antidepressant to the site of discomfort, for example by injection, supersonic powder injection, transdermal electroporation, topical application, or encapsulation in a slow release delivery vehicle.

15

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a graph illustrating the difference in the threshold pain response over 22 days in the ipsilateral and contralateral paws of rats subjected to nerve
20 ligation. The row of bars at the back of the figure illustrates the static mechanical allodynia 50% withdrawal threshold (grams) of nerve-ligated rats, and the row of bars at the front of the figure illustrates the thermal hyperalgesia (seconds) on successive days following spinal nerve ligation. Values depict group means (n=6 per group).

25 **Figures 2A-C** are a series of graphs illustrating the time course of the effect of systemic amitriptyline (AMI) at various doses on thermal hyperalgesia in the ipsilateral paw of nerve-ligated rats (Figure 2A), and the thermal withdrawal threshold in the contralateral paw (Figure 2B), with ●=saline control; ■=1mg/kg AMIT; ▲=3 mg/kg AMI, and ▼=10 mg/kg AMI. Figure 2C illustrates the cumulative change in

the Maximum Possible Effect (Δ MPE) time course for ipsilateral (ipsi) and contralateral (contra) paws, with open bars representing saline control, bars with rightward sloping lines representing 1 mg/kg AMIT, bars with leftward sloping lines representing 3 mg/kg AMI, and bars with crosshatching representing 10 mg/kg AMI.

- 5 Values represent the MPE \pm s.e.m. (n=5 for 1 and 3 mg/kg, and n=15 for 10 mg/kg).
 *=P<.05 compared to saline controls.

Figures 3A-C are a series of graphs illustrating the time course of the effect of spinal amitriptyline (60 μ g) on thermal hyperalgesia in the ipsilateral paw (Figure 3A), and the thermal withdrawal threshold in the contralateral paw (Figure 3B), with 10 3A), and the thermal withdrawal threshold in the contralateral paw (Figure 3B), with \bullet =saline control, \blacksquare = 60 μ g AMIT. Figure 3C represents the cumulative change in the MPE (Δ MPE) time course in ipsilateral (ipsi) and contralateral (contra) paws, with open bars representing saline control, rightward sloping bars representing 100 nmol AMI (ipsi), and leftward sloping bars representing 100 nmol AMI (contra). Values 15 represent the MPE \pm s.e.m. (n=6 for both groups). *=P<.05 compared to saline controls.

Figures 4A-C are a series of graphs illustrating the time course of the effect of local injection of 100 nmol amitriptyline on thermal hyperalgesia. Figure 4A shows 20 the effect in the ipsilateral paw when the antidepressant is injected into the nerve-injured paw (ipsi) and non-injured paw (contra), and Figure 4B shows the time course of the change in the thermal withdrawal latency in the contralateral paw, when the antidepressant is injected into the nerve injured paw (ipsi) and non-injured paw (contra), with \bullet =saline control; \blacksquare =AMI ipsi, and \blacktriangle = AMI contra. Figure 4C 25 depicts the cumulative change in MPE value (Δ MPE), with open bars representing saline control (ipsi), bars with rightward sloping lines representing 100 nmol AMI (ipsi), and bars with leftward sloping lines representing 100 nmol AMI (contra). Values represent the MPE \pm s.e.m. (n=9 for all groups). *=P<.05 as compared to saline controls, t =P<.01 as compared to contralateral controls.

Figures 5A-C are a series of graphs illustrating the time course of the effect of systemic amitriptyline on static mechanical allodynia in the ipsilateral paw (Figure 5A) and on the threshold of withdrawal to static mechanical stimuli in the contralateral paw (Figure 5B), with ●=saline control; ■=1.5 mg/kg AMI; ▲=5 mg/kg AMI, and ▼=10 mg/kg AMI. Figure 5C represents the cumulative change in the MPE (Δ MPE), with open bars representing saline control, bars with rightward sloping lines representing 1.5 mg/kg AMI, bars with leftward sloping lines representing 5 mg/kg AMI, and bars with crosshatching representing 10 mg/kg AMI. Values represent MPE \pm s.e.m. (n=5 for 1.5 and 5 mg/kg, and n=15 for 10 mg/kg). *= P <.05 as compared to saline, t = P <.05 as compared 1.5 and 5.0 mg/kg.

Figures 6A-C are a series of graphs illustrating the time course of the effect of spinal amitriptyline on static mechanical allodynia in the ipsilateral paw (Figure 6A), on the threshold of withdrawal to static mechanical stimuli in the contralateral paw (Figure 6B), with ●=saline control; ■=30 μ g AMI, ▲=60 μ g AMI, and ▼=90 μ g AMI. Figure 6C illustrates the cumulative change in the MPE (Δ MPE), with open bars representing saline control; bars with rightward sloping lines representing 30 μ g AMI, bars with leftward sloping lines representing 60 μ g AMI, and bars with crosshatching representing 90 μ g AMI. Values represent the MPE \pm s.e.m. (n=3 for 30 μ g and 90 μ g, and n=5 for 60 μ g). *= P <.05 compared to saline controls.

Figures 7A-C are a series of graphs illustrating the time course of the effect of local peripheral injection of amitriptyline (100nmol) on mechanical allodynia. Figure 7A illustrates the effect in the ipsilateral paw when the antidepressant is injected locally into the nerve-injured paw (ipsi) and non injured paw (contra). Figure 7B illustrates the time course of the change in the mechanical threshold of the contralateral paw following injection into the nerve-injured paw (ipsi) and the non-injured paw (contra), with ●=saline control; ■=AMI (ipsi), and ▲= AMI (contra).

Figure 7C illustrates the cumulative change in the value of MPE (Δ MPE) over the entire time course), with open bars representing saline control (ipsi), bars with rightward sloping lines representing 100 nmol AMI (ipsi), and bars with leftward sloping lines representing 100 nmol AMI (contra). Values represent the MPE \pm s.e.m. (n=9 for all groups). *= $P < 0.05$ as compared to saline controls, and $t = P < 0.01$ as compared to contralateral controls.

Figures 8A-B are two graphs illustrating the time course of the local antinociceptive effect produced by coadministration to rats of amitriptyline (AMI) with formalin 2.5% as measured by number of flinches (Figure 8A) or time spent biting or licking (Figure 8B). ●=2.5% formalin alone; open triangle = 2.5% formalin+10nM AMI; open square=2.5% formalin+30 nmol AMI; open diamond = 2.5% formalin+100 nmol AMI. Values depict group means (n=5-11 per group); error bars for s.e.m. were omitted in the interest of clarity.

Figures 9A-C are a series of bar graphs illustrating the cumulative scores for phase 1 flinches (Figure 2A), phase 2 flinches (Figure 9B), and biting/licking time (Figure 9C) following coadministration to rats of amitriptyline (100 nmol) with 2.5% formalin (F). The antinociceptive effect of amitriptyline administered to the ipsilateral paw (local effect) was not reproduced by administration of amitriptyline into the contralateral (CONT) hindpaw. The values depicted are \pm s.e.m. values (n=5-11). *= $P < 0.05$, **= $P < 0.01$, and ***= $P < 0.001$ compared to the formalin alone group.

Figures 10A-C are a series of graphs illustrating the inhibitory effect on phase 1 flinches (Figure 10A), phase 2 flinches (Figure 10B), and time spent biting and licking (Figure 10C) following coadministration to rats of amitriptyline (100 nmol) (open diamonds) on formalin-induced behaviors at different concentrations of formalin (filled circles). n=6-11 per group. *= $P < 0.05$, **= $P < 0.01$, and ***= $P < 0.001$ compared to the formalin only group.

Figures 11A-C are a series of graphs illustrating partial reversal of the local antinociceptive action of amitriptyline in the 2.5% formalin (F) test by coadministration with caffeine (CAFF) as measured by phase 1 flinches (Figure 11A), phase 2 flinches (Figure 11B), and time spent licking and biting (Figure 11C). n=10-15 except for the caffeine (1500 nmol) group where n=5. *= $P < 0.05$, ***= $P < 0.001$ compared to the formalin only group; t = $P < 0.05$, tt = $P < 0.01$ compared to formalin/amitriptyline group. The filled bars illustrate 2.5% formalin alone, the empty bars illustrate administration of AMI alone, the bars with leftward sloping lines illustrate AMI+500 nmol CAFF, and the crosshatched bars illustrate 1500 nmol caffeine alone or with 100 nmol AMI.

Figures 12A-B are two graphs illustrating the time course of caffeine (CAFF) reversal of amitriptyline (AMI) action in the 2.5% formalin (F) test for the experiment illustrated in Figures 4A-C as measured by number of flinches (Figure 12A) and time spent biting and licking (Figure 12B). ●=2.5% formalin alone; open upward-pointing triangle=2.5% F+100 nmol AMIT; ■=2.5% F+100 nmol AMI+1500 nmol CAFF; ♦=2.5% F+1500 nmol CAFF. Values depict group means (n=5-11 per group); error bars for s.e.m. were omitted in the interest of clarity.

Figures 13A-B are graphs illustrating the reversal of the antinociceptive action of amitriptyline (AMI) against 1.5% formalin (F) by caffeine (CAFF) (Figure 13A), and 8-cyclopentyl-1,3-dimethylxanthine (CPT) (Figure 13B). n=5-6 per group. **= $P < 0.01$, ***= $P < 0.001$ compared to formalin alone; t = $P < 0.05$, ttt = $P < 0.001$ compared to formalin/amitriptyline. The filled bars represent the effect of administering 1.5% formalin alone; the bars with downward sloping lines represent CAFF alone or in combination with AMIT, and the open bars represent the effect of AMI alone.

Figure 14 is a bar graph illustrating enhancement of the local antinociceptive of 30 nM and 100 nM amitriptyline (open bars) against 2.5% formalin (filled bar) action as measured by decrease in phase 2 flinches in the formalin test by coadministration of the adenosine kinase inhibitor NH_2dAD (100nmol) (bars with rightward sloping lines), but not by the adenosine deaminase inhibitor 2'-deoxycytosine (DCF) (100 nmol) (bars with cross hatching). n=6-8 per group. $\ast=\text{P}<0.05$ compared to formalin alone; $\text{t}=\text{P}<0.05$ compared to the formalin/amitriptyline combination.

Figures 15A-B are bar graphs illustrating inhibition of responses by rats to 2.5% formalin (filled bars) when the formalin is coadministered with MK-801 (20 nM or 100 nM) (open bars) into the ipsilateral (I) paw, or the contralateral (C) paw. n=6-8 per group. $\ast=\text{P}<0.05$; $\ast\ast=\text{P}<0.01$, and $\ast\ast\ast=\text{P}<0.001$ compared to formalin alone.

Figure 16A-B are bar graphs illustrating that the antinociceptive action of amitriptyline (20nmol and 100nmol) against 2.5% formalin (F) (dark bar) was not decreased by coadministration of phentolamine (PTA) (100nmol and 500nmol) (bars with crossed hatching) or naloxone (NLX) (100nmol and 300 nmol) (bars with horizontal lines). n=5-8 per group. $\ast=\text{P}<0.05$, $\ast\ast\ast=\text{P}<0.001$ compared to the formalin group; $\text{t}=\text{P}<0.05$ compared to the formalin/amitriptyline combination.

Figures 17A-C are a series of graphs showing dose-related inhibition of formalin-induced pain response behaviors by locally administered tricyclic antidepressants. The bar graphs depict the control values for formalin (solid bar) and saline (open bar) when injected contralaterally; while the dosage graphs depict the results when antidepressant, rather than control, is injected. Figures 17A and 17B show amelioration of phase 1 and phase 2 flinching behaviors, respectively, and

Figure 17C shows amelioration of biting/licking behaviors induced by formalin 2.5% (■) injected into the dorsal hindpaw of the rat when the formalin is coadministered with desipramine (DES) or fluoxetine (FLUOX) into the dorsal hindpaw (local effect) or injected into the contralateral hindpaw (systemic effect). ▲=ipsilateral DES; open triangle=contralateral DES; ●=ipsilateral FLUOX; ○=contralateral FLUOX. Data depicts mean \pm s.e.m. for n=5-6; n=8 for ipsilateral control). *= $P<0.05$, **= $P<0.01$, ***= $P<0.001$ compared to corresponding formalin group.

Figures 18A-D are a series of graphs illustrating the suppression of pain related behaviors by coadministration of desipramine with 2.5% formalin. Figures 18A and 18B show the time course for the suppression of flinching behaviors by coadministration of DES with the formalin into the ipsilateral (Figure 18A) or contralateral (Figure 18B) paw. Figures 18C and 18D show the time course for suppression of biting/licking behaviors after administration of the highest doses injected into the ipsilateral (Figure 18C) or contralateral (Figure 18D) paw. Values are means, with error bars omitted in the interest of clarity. Data corresponds to Figure 17.

Figures 19A-C are a series of graphs illustrating the results of local injection of desipramine (DES) (100 nmol) and fluoxetine (FLX) (100 nmol). Figure 19A depicts the local antinociceptive action of desipramine, but not fluoxetine, when injected into the ipsilateral (ipsi) paw corresponding to the spinal nerve ligation, and Figure 19B depicts the lack of effect in the contralateral paw (contra) with ●=saline control; ■=DES; and ▲=FLUOX. Figure 19C shows the cumulative change in MPE with the open bars representing saline control, the rightward sloping bars representing DES and the leftward sloping bars representing FLUOX. n= 9 per group. *= $P<0.05$ compared to saline group.

Figures 20A-B are two graphs illustrating the effect of caffeine (1500 nmol) (CAFF) for inhibiting the antinociceptive effect of desipramine when coadministered into the paw ipsilateral to spinal ligation. Figure 20A shows the anti-hyperalgesia MPE over time wherein ●=100nmol DES; ■= 1500 nmol CAFF; and ▲= DES + CAFF. Figure 20B shows the cumulative change in MPE (Δ MPE) wherein the open bar represents 100nmol DES; the bar with rightward sloping lines represents 1500 nmol CAFF; and the bar with crosshatching represents DES + CAFF. n=9 per group; *= $P < 0.05$ compared to saline group.

Figures 21A-C are a series of graphs showing a lack of suppression of the local antinociceptive effect of desipramine (DES) or fluoxetine (FLUOX) caused by coadministration of caffeine (CAFF) (1500 nmol) in the 1.5% formalin test. Figure 21A shows the effect on phase 1 flinches, and Figure 21B shows the effect on phase 2 flinches. Figure 21C shows the effect phase 2 biting/licking time. The empty bar represents the saline control; the bar with rightward sloping lines represents the effect of caffeine (CAFF) either alone or in combination with the test drug. Values depict mean \pm s.e.m. for n=6 per group. *= $P < 0.05$, **= $P < 0.01$, ***= $P < 0.001$ compared to formalin group. NS indicates no significant difference from corresponding antidepressant group.

20

Figures 22A-B are graphs illustrating the effect of locally administered desipramine and fluoxetine on paw volume in the presence of formalin. Figure 22A shows the effect of saline (control), desipramine, or fluoxetine on paw volume following coadministration with formalin into the ipsilateral paw in the presence of formalin. ●=formalin+saline; open diamond=formalin+100 nmol FLUOX; open downward pointing triangle=formalin+300 nmol DES; and open square=formalin+300 nmol FLUOX. Figure 22B shows the effect on paw swelling of desipramine or fluoxetine injected at either concentration used in Figure 22A into the contralateral hindpaw (systemic effect). Paw volumes were determined at the end of

the behavioral observation period. n=5-6 per group. ***=P<0.001 compared to the saline-treated group.

Figures 23A-C are a series of graphs illustrating the effects of desipramine, fluoxetine, and amitriptyline as compared with saline control on paw volume when injected into the dorsal hindpaw of the rat. Figure 23A shows the effects of 100nmol DES (open diamond) and 300nmol DES (open square). Figure 23B shows the effects of 10nmol FLUOX (open upward facing triangle); 30nmol FLUOX (open downward-facing triangle), 100nmol FLUOX (open diamond), and 300nmol FLUOX (open square). Figure 23C shows the effects of 100nmol (open triangle) and 300nmol (open square) amitriptyline on dorsal paw volume. n=6 per group. *=P<0.05, ***=P<0.001 compared to the saline-injected group.

Figures 24A-B are graphs illustrating the effects of the selective 5-HT receptor antagonists propranolol (5-HT₁), ketanserin (5-HT₂), tropisetron (5-HT_{3/4}) and GR113808A (5-HT₄) on the increase in paw volume induced by local injection of fluoxetine (FLX) (100 nmol). Figure 24A shows the percent increase in paw volume over a three hour period of 100nmol FLX alone (●); FLX + 300nmol propranolol (open diamond); FLX + 500nmol ketanserin (□), FLX + 300nmol tropisetron (open upward-pointing triangle); and FLX + 500nmol GR113808A (open downward-pointing triangle). Figure 24B shows the cumulative effect of the increase of Figure 24A over the same time period wherein the open bar represents FLX alone; the bar with horizontal lines illustrates FLX+propranolol; the bar with vertical lines illustrates FLX+ketanserin; the bar with rightward sloping lines represents FLX+tropisetron; and the bar with leftward sloping lines represents FLX+GLR113808A. n=5 per group. *=P<0.05, **=P<0.01, ***=P<0.001 compared to the fluoxetine group.

Figure 25 is a graph showing the lack of effect at dosages of 50 nmol, 150 nmol and 500 nmol ketanserin (KET), a selective 5-HT receptor antagonist, on

formalin-evoked behaviors (phase 2 flinches) in the rat 1% formalin test. Values depict the mean of 6 measurements per group. ● = formalin 2.5% control; □ = propranolol;

Δ = ketanserin; open diamond = tropisetron; open hexagon = GR 113808A.

5 * = P<0.05 compared to formalin controls.

Figure 26 is a graph showing the lack of effect of ketanserin (KET) at dosages of 50 nmol and 500 nmol on formalin-evoked behaviors (flinches) in the rat 1% formalin test. Values depict the mean of 6 measurements per group. ■ = 1% formalin control; □ = 50 nmol KET; ▲ = 500 nmol KET.

Figures 27A and B are graphs showing blockade of rat paw edema induced by local injection of 3 nmol 5-HT by coadministration of 3 nmol or 30 nmol ketanserin with 5-HT. Figure 27A is a graph illustrating the cumulative increase in paw volume over 180 minutes. Open bars = 5-HT alone; cross-hatched bars = 5-HT + KET (3 nmol 5-HT = left hand columns; 30 nmol 5-HT = right hand columns). Figure 27B is a graph illustrating the time course of action in terms of % increase in paw volume over 180 minutes. ■ = 5-HT 3 nmol alone; □ = 5-HT 3 nmol + KET 500 nmol; ● = 5-HT 30 nmol alone; ○ = 5-HT 30 nmol + KET 500 nmol.

20 ** = P<0.01, *** = P<0.001 compared to 5-HT response.

Figures 28A-C are a series of graphs illustrating the effects of atropine, mecamylamine and d-tubocurarine at dosages of 10, 30, 100, 300 and 1000 nmol on flinching and biting/licking behaviors produced by 2.5% formalin in rats. Figure 28A illustrates the effects of the drugs on the number of phase 1 flinches. Figure 28B illustrates the effects of the drugs on the number of phase 2 flinches, and Figure 28C illustrates the effects of the drugs on phase 2 biting/licking time. (dark hexagon = formalin 2.5%; ○ = amitriptyline; □ = mecamylamine; Δ = atropine (ipsilateral); dark

triangle = atropine (contralateral); open diamond = d-tubocurarine.

* = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$ compared to respective formalin controls.

DETAILED DESCRIPTION OF THE INVENTION

5

In accordance with the present invention, a new method has been discovered for producing local analgesia by local administration of a tricyclic, second generation or third generation antidepressant to a body site of local discomfort.

10 Examples of tricyclic antidepressants useful in the practice of this invention are those having the structure:





wherein Z is a 7-membered ring, optionally containing 1 or 2 biocompatible heteroatoms, or an 8-membered bicyclic ring,

15 Ar_1 and Ar_2 are optionally substituted aromatic rings fused to Z, and

R is an alkylamino or arylamino substituent.

Examples of Z as an 8-membered bicyclic ring include (2,2,2,)bicyclooctane and (3,2,1) bicyclooctane, and the like. Oxygen and nitrogen are the preferred heteroatoms for optional inclusion in Z, the central ring of the three ring structure, but
 20 sulfur can also be included. R can be an alkylamino or arylamino substituent, or an N-oxide derivative thereof and generally comprises from 4 to 5 carbons, although as few as two carbons or as many as seven carbons, or even as many as 10 carbons can be contained in R. If R is an alkylamino substituent, in some embodiments the alkylamino is a tertiary or secondary amino group. For example, R can be





The preferred point of attachment of R to the tricyclic structure is at carbon 5, or to a heteroatom located at position 5 in the ring structure.

5

Examples of the optional biocompatible halogen substituent for Ar_1 and/or Ar_2 are chlorine, bromine and fluorine, with chlorine being preferred. For example Ar_1 and/or Ar_2 can be substituted with one or two of the biocompatible halogen atoms.


10

Non-limiting examples of tricyclic antidepressants useful in the practice of the present invention are clomipramine, imipramine, amitriptyline, desipramine, nortriptyline, amoxapine, maprotiline, trimipramine, and the like, and suitable combinations of any two or more thereof. Amitriptyline and desipramine are presently preferred.

15

Additional antidepressants contemplated for use herein include compounds sometimes referred to as "second-generation" or "third-generation" antidepressants, having pharmacokinetics and potency similar to those of the tricyclic antidepressants described herein. Examples of second generation or third generation antidepressants useful in the practice of the invention are those having the structure

20



wherein Ar_3 is an N-containing heterocyclic ring, e.g., a substituted 1,2,4-triazole or triazone,

Y is either an aryl group fused to the heterocyclic ring or one or two substituents selected from alkyl, alkyloxy, arylalkyl, arylalkyloxy, aryl and heteroaryl substituents comprising a total of about 4 to 8 carbons attached to Ar_3 ,

25

X is an alkyl group comprising 2 to 5 carbon atoms linking Ar_3 and Ar_4 ,

Ar_4 is a piperazine attached to X by a first nitrogen atom of Ar_4 , and

Q is benzene ring optionally substituted with a biocompatible halogen, and attached to Q at a second nitrogen atom of Ar₄.

For example, X can contain 3 carbons, and Ar₃ can be a 1,2,4-triazone substituted at the 4 position with an arylalkyloxy substituent containing 6 to 8 carbon atoms. In an embodiment presently preferred, Y is a heteroarylalkyl substituent containing an oxygen atom.

The preferred biocompatible halogen substituents for Q are selected from chlorine, bromine and fluorine.

Representative examples of second generation or third generation antidepressants useful in the practice of the present invention are trazodone, bupropion, mirtazapine, venlafaxine, fefazodone, and the like.

15

In one aspect, the present invention includes compositions for local administration of tricyclic, second generation or third generation antidepressants to a subject having a site of local discomfort. Such compositions are specifically formulated to ensure substantially local, rather than systemic administration of such antidepressant(s). In addition to one or more of the antidepressants described for use herein, the compositions of the invention may comprise one or more adjuvants which facilitate delivery, such as inert carriers, penetration enhancing agents, colloidal dispersion systems.

Representative and non-limiting examples of such inert carriers can be selected from water, isopropyl alcohol, gaseous fluorocarbons, ethyl alcohol, polyvinyl pyrrolidone, propylene glycol, a fragrance, a gel-producing material, stearyl alcohol, stearic acid, spermaceti, sorbitan monooleate, methylcellulose, and suitable combinations of two or more thereof.

The topical formulation can also include one or more known penetration enhancing agents selected to have minimal allergic or irritating effect. Known penetration enhancing agents include DMSO and azacyclo compounds, such as those disclosed in U.S. Patent Nos. 4,755,535, 4,801,586, 4,808,414, and 4,920,101, which are incorporated herein, each in its entirety. No more of a penetration enhancing agent should be included in such compositions than is consistent with the goal of attaining substantially local delivery, rather than systemic delivery, of the tricyclic, second generation or third generation antidepressant(s). Those of skill in the art will be able to select a suitable penetration enhancing agent useful for the particular degree of penetration desired.

In another embodiment, the invention composition is formulated for local injection. In such an embodiment, the composition generally comprises a physiologically compatible saline solution and may optionally be encapsulated in a slow release delivery vehicle suitable for local injection, such as a colloidal dispersion system or in polymer stabilized crystals. Colloidal dispersion systems include nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome or microsphere. Liposomes are artificial membrane vesicles which are useful as slow release delivery vehicles when injected or implanted, or when contained within a topical preparation. One type of microspheres suitable for encapsulating drugs for local injection is poly(D,L)lactide microspheres, as described in D. Fletcher, *Anesth. Analg.* 84:90-94, 1997.

The invention compositions may be formulated for topical administration, for instance as a cream, a lotion, a gel, an ointment, a spray, or an aerosol. Tricyclic, second generation and third generation antidepressants are quite lipid soluble themselves and, readily cross lipid barriers to gain access to central nervous system

sites. This same property enables them to cross the perineurial barrier surrounding peripheral nerve endings (this barrier limits access of some agents to such sites, I. Antonijevic et al., Perineurial defect and peripheral opioid analgesia in inflammation, *J. Neurosci* 15 (1995) 165-172), giving them ready access to the peripheral nerve terminal and adjacent sites. Topical administration can also be effected by delivering the invention compositions via a slow release "patch." Therefore, in yet another embodiment, the composition further includes a patch delivery system having the antidepressant sequestered for slow release upon application to skin or mucous membrane.

10

In another aspect of the invention, there are provided methods for producing local analgesia in a subject, such as a mammal, by locally administering an effective amount of a tricyclic, second generation or third generation antidepressant(s) to a site of local discomfort. The term "local administration" includes any method of local administration known in the art. The antidepressant(s) used in the practice of the invention can be injected locally, for example, by subcutaneous or intra-articular injection. Alternatively, the tricyclic, second generation or third generation antidepressant(s) can be applied topically by applying a cream ointment, spray, or gel comprising the any antidepressant contemplated for use herein to an area of skin adjacent to a site of pain. Additional routes of local administration contemplated herein include topical application, local injection or subcutaneous implantation of the antidepressant encapsulated in a slow release vehicle, such as a liposome or microsphere.

15

20

25

The term "effective amount" means the quantity of a compound according to the invention necessary to prevent, to cure, or at least partially arrest a symptom of local pain or discomfort in a subject. A subject is any mammal, preferably a human. Amounts effective for creating a substantially local therapeutic effect will, of course, depend on the severity of the disease causing the painful condition, and the weight

and general state of the subject. Typically, animal models, such as those described in the Background and Examples herein, may be used to determine suitable dosages to be used to achieve substantially local delivery of the tricyclic, second generation or third generation antidepressant(s). A recent pain scale developed by Galer et al.

- 5 (Development and preliminary validation of a pain measure specific to neuropathic pain: The Neuropathic Pain Scale, *Neurology* 48 (1997) 332-338), which uses terminology specific for neuropathic pain, should be better able to delineate the symptoms within the syndrome. In addition, various general considerations taken into account in determining the "therapeutically effective amount" are known to those
- 10 of skill in the art and are described, *e.g.*, in Gilman *et al.*, eds., Goodman And Gilman's: The Pharmacological Bases of Therapeutics, 8th ed., Pergamon Press, 1990; and Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Co., Easton, Pa., 1990, each of which is herein incorporated by reference.

- 15 The compositions and methods of the invention are particularly suited to producing local analgesia for painful symptoms known in the art as "neuropathic pain." As used herein the term "neuropathic pain" refers to pain syndromes known to be neuropathic (*i.e.*, due to lesions or dysfunction in the nervous system) including certain relatively generalized syndromes, such as peripheral neuropathy, phantom
- 20 pain, reflex sympathetic dystrophy, causalgia, central pain, syringomyelia, painful scar, and the like. Certain relatively localized syndromes are also considered to be neuropathic. Among these are specific neuralgias at any location of the body, head or face; diabetic, alcoholic, metabolic or inflammatory neuropathies; post herpetic neuralgias; post traumatic and post endodontic odontalgia; thoracic outlet syndrome;
- 25 cervical, thoracic, or lumbar radiculopathies with nerve compression; cancer with nerve invasion; post traumatic avulsion injuries; post mastectomy pain, post thoracotomy pain; post spinal cord injury pain; post stroke pain; abdominal cutaneous nerve entrapments; primary tumors of neural tissues; and arachnoiditis, and the like.

Other pain syndromes believed to have a neuropathic component are stump pain, fibromyalgia, regional sprains or strains (crushing injury), myofascial pain, psoriatic arthropathy, polyarteritis nodosa, osteomyelitis, burns involving nerve damage, AIDS related pain syndromes, and connective tissue disorders, such as
5 systemic lupus erythematosus, systemic sclerosis, polymyositis, and dermatomyositis, and the like.

The compositions and methods of the invention are also particularly suited to producing local analgesia for symptoms of localized pain and discomfort caused by
10 inflammatory conditions. Inflammatory conditions that can be treated in the practice of the invention include conditions of acute inflammation (e.g. trauma, surgery and infection) or chronic inflammation (e.g., arthritis and gout).

The relative high concentrations of drugs attainable by local administration,
15 coupled with a lesser incidence of the side effects characteristic of systemic absorption, produce particular benefits in treatment of discomfort associated with inflammatory or neuropathic pain conditions using the compositions and methods of this invention.

20 A rat spinal nerve ligation (SNL) model was used to determine the effect of tricyclic, second generation or third generation antidepressant(s) against neuropathic pain as described in full in the Examples herein. In this model, locally administered amitriptyline and desipramine, tricyclic antidepressants, were shown to be effective in alleviating thermal hyperalgesia. The local anti-hyperalgesic effects of amitriptyline
25 and desipramine administered to a nerve-damaged hindpaw were observed without a concomitant analgesic effect on the contralateral paw. The maximal effect of locally administered amitriptyline was equal in magnitude to that of systemic injection, but alleviated manifestations of neuropathic pain over a longer time period than did systemic administration of the drug. This local effect produced no change in the

thermal threshold of the contralateral paw or the paw of naive animals following local administration. Further, the local nature of this effect was verified when injections into the contralateral paw were found to have no significant effect on ipsilateral response latencies.

5

In contrast to the beneficial actions of locally administered amitriptyline against thermal hyperalgesia, amitriptyline was observed to be ineffective in the treatment of nerve injury-induced mechanical allodynia. There may even be a slight untoward effect following spinal administration of amitriptyline (60 μ g) that became statistically significant when the cumulative change in the maximum possible effect (Δ MPE) was analyzed (Figure 6C). Increasing the spinal dose of amitriptyline to 90 μ g, however, caused a significant sedative effect so that the animals were flaccid in the testing apparatus, a result that hindered their ability to respond to the stimuli. An interesting observation with the use of amitriptyline in the allodynia test was the decrease in response threshold observed in the contralateral paw (Figs. 5A-C, 6A-C, and Figures 7A-C), an effect referred to herein as a "hyperaesthetic tactile response." While the hyperaesthetic response threshold was similar to that of the nerve injured paw qualitatively, the observed behavioral response (a brisk withdrawal) differed from that exhibited by a normal neuropathic pain response (i.e., biting, licking, and guarding). The results of this study indicate that amitriptyline preferentially alleviates thermal hyperalgesia through all routes of administration, but has no effect on allodynia.

These findings suggest that the pathophysiology of thermal hyperalgesia and tactile allodynia are maintained through distinct mechanisms, which has also been suggested by the differential profile of other drugs like dextromethorphan on these two endpoints (M. Tal and G.J.Bennett (1994) *Neuroreport* 5:1438-1440). Therefore, the efficacy of tricyclic antidepressants, such as amitriptyline in a patient may be determined by which of the neuropathic symptoms is the most strongly expressed.

For example, in a patient presenting with robust allodynic like symptoms, amitriptyline and antidepressants with similar activity may not make any perceived difference in the pain. However, if the dominant complaint is one that can be characterized as hyperalgesia, amitriptyline and drugs with similar activity may provide relief.

The experiments upon which this invention is based also demonstrate a local peripheral antinociceptive action for antidepressants, such as tricyclic, second generation and third generation antidepressants, against inflammation. In the rat formalin test, a well known animal model for inflammatory conditions, the antinociceptive action is shown by reduction in pain responses (both flinching behaviors and biting/licking time) caused by injection of formalin into a rat paw. The local nature of this action is verified in the animal model by the lack of effect of amitriptyline injected into the contralateral hindpaw. The antinociceptive effect is seen at locally injected doses of amitriptyline up to 100 nmol, which corresponds to a systemic dose of 0.22 mg/kg (for a 142 g rat). In the 2.5% formalin test, amitriptyline, desipramine and fluoxetine 10-300 nmol produced a dose-related reduction in phase 2 (16-60 min) flinching and in biting/licking behaviors when coadministered with the inflammatory irritant, formalin. Phase 1 flinch behaviors (0-12 min) were reduced at the highest dose. This action was locally mediated, as it was not seen when the antidepressants (100, 300 nmol) were injected into the contralateral hindpaw. The local action of desipramine and fluoxetine was partially altered by coadministration of caffeine 1500 nmol, but the action of amitriptyline was reversed by caffeine.

25

The peripheral action of amitriptyline involves an interaction with endogenous adenosine systems as shown by the Examples herein. A similar peripheral antinociceptive action of NH_2dAD , an inhibitor of adenosine kinase, which also acts

by changing the extracellular availability of adenosine, is shown in the rat formalin test.

5 The local effect of locally administered tricyclic, second generation or third generation antidepressants as demonstrated in the formalin test is quite robust, with phase 1 and phase 2 flinching behaviors and the total biting/licking time being suppressed markedly.

10 While the mechanism of action of the antidepressants does not form a part of this invention, it has been determined that the peripheral antinociceptive action of amitriptyline involves endogenous adenosine with a subsequent activation of adenosine A₁ receptors on sensory nerve terminals (reviewed Sawynok J. (1997) In: *Purinergic Approaches to Experimental Therapeutics* (Ed. Jacobsen K., Jarvis M.) Wiley-Liss Inc., New York, 495) This observation is based on the ability of caffeine, 15 a non-selective adenosine A₁ and A₂ receptor antagonist, and CPT, a selective adenosine A₁ receptor antagonist, to inhibit the action of amitriptyline on phase 2 flinching responses and biting/licking time. Phase 1 responses were unaffected by caffeine, and may result from another mechanism.

20 The studies upon which this invention is based also demonstrate that an inhibitor of adenosine kinase (NH₂dAD) can augment the local action of amitriptyline. Adenosine kinase is an intracellular enzyme (reviewed J.D.Geiger et al., *Regulators of endogenous adenosine levels as therapeutic agents in: Purinergic approaches in Experimental Therapeutics*, Ed. Jacobsen, K.A. and Jarvis, M.F., Wiley-Liss Inc., 25 New York (1997) 55-84) and its inhibition would result in an intracellular accumulation of adenosine and an efflux of adenosine from the cell along a concentration gradient. This effect would enhance the extracellular adenosine available for uptake and the interaction with amitriptyline. Inhibition of adenosine

deaminase with 2'-deoxycoformycin did not produce the same response, and this is generally consistent with the relative intrinsic effects expressed by these two agents.

It is important to note that in the experiments described herein all of these influences on adenosine availability are superimposed on an inflammatory stimulus (formalin). Endogenous adenosine is released peripherally by formalin (Doak GJ, Sawynok J. (1995) *Eur J Pharmacol* 281: 311), as well as by inflammatory processes in general (Cronstein, 1994). Thus, the stimulus generates the substrate (adenosine and its uptake) with which amitriptyline interacts. The lack of, or more limited, effect of amitriptyline observed at lower concentrations of formalin inflammatory model (Figures 10A-C) may well be due to an inadequate release of endogenous adenosine at the lower stimulus intensity.

Similarly, the experiments on which this invention is based have shown that caffeine and CPT were without an intrinsic effect at 1.5% formalin, or slightly reduced the effect of formalin at 2.5% (cf. Figures 11A-C, 12A-B and 13A-B). The lack of effect of CPT indicates that the amount of endogenous adenosine generated by the formalin was insufficient to produce antinociception; such a response only occurred when amitriptyline was present and inhibited adenosine uptake. The modest effect of caffeine at 2.5% formalin may reflect a slight predominance of adenosine A₂ receptor activation in this experiment. Caffeine has been shown to lack a peripheral intrinsic effect due to blockade of opposing influences of adenosine A₁ and A₂ receptors (Doak G.J., Sawynok J. (1995, *supra*).

In the spinal nerve ligation model, amitriptyline (100 nmol) and desipramine (100 nmol), but not fluoxetine (100 nmol), produced a local and complete antihyperalgesic action in the hindpaw corresponding to the ligated side when thresholds were determined using a thermal paw stimulator. Desipramine and amitriptyline, at 100 and 300 nmol, which are maximally effective doses, produced a

slight increase in paw volume, but fluoxetine produced a robust and sustained increase in paw volume over the complete dosage range (10-300 nmol). No agent significantly altered the degree of paw swelling produced by formalin in the inflammatory pain model when coinjected with the formalin. The increase in paw volume produced by fluoxetine was inhibited by ketanserin, a selective 5-HT₂ receptor antagonist, but not by other selective 5-HT receptor antagonists. The pronounced local pain alleviating actions in the absence of marked changes in paw volume produced by the tricyclic antidepressants desipramine and amitriptyline in the formalin test and the spinal nerve ligation model indicate that tricyclic, second generation or third generation antidepressant(s) can be used to produce local analgesia, for example in inflammatory and neuropathic pain states.

The invention will now be described in greater detail by reference to the following non-limiting examples.

EXAMPLE 1

Experiments were conducted on male Sprague-Dawley rats (100-120 gm) (Charles River, Quebec, Canada). Animals were housed in pairs, maintained on a 12:12 hour light-dark cycle at 22°±1°C and given ad-libitum access to food and water.

A. Preparation of the spinal nerve ligation animal model

Surgery on male rats was performed using aseptic technique under a dissection microscope and with adherence to the guidelines of the IASP on animal experimentation in pain research. The neuropathic condition was created by a tight ligation of the dorsal roots of lumbar nerves L-5 and L-6. Sprague Dawley rats (100-125 gm) were anaesthetized and surgically maintained using Halothane (1.5-2.0%). The animals were surgically clipped in the dorsal pelvic area. Antibacterial eye drops

were applied to the eyes to prevent drying through the course of the surgical procedure. The animals were placed on a water heated pad ($37\pm0.5^{\circ}\text{C}$). The area for surgical incision was cleansed and sterilized with ethyl alcohol and iodine. A 3 cm incision was made along the dorsal midline with the position of the iliac crest indicating the incision midline. A stab incision was made on the left side of the vertebral column at the level of the sacrum. The ligament attachments to the spinal cord were cut 2 cm rostral to the stab incision. Blunt dissection was used to further disconnect and separate the remaining tissue from the spinal cord. One retractor was placed at the L-4 vertebrae and a second retractor was used to counter retract the muscle away from the sacro-iliac crest. Blunt dissection was used to clear the sacro-iliac crest, the L-6/S-1 articular (facet) joint, and the transverse process of L-6. The facet was nipped off with rongeurs. A third retractor was used to retract the muscle away from the L-6 transverse process.

Under the dissecting microscope and using jewelers forceps, the connective tissue between the L-6 transverse process and the sacro-iliac rim was removed. Rongeurs were used to slowly nip away the transverse process of L-6 and expose the area for access to the L-5/L-6 nerve bundle. Using a glass probe the L-4/L-5 nerve bundle was hooked and manipulated to isolate the L-5 nerve. Once free of the L-4 nerve, the L-5 nerve was tightly ligated using 6-0 silk so that the nerve bulged on both sides of the ligature. The suture ends were trimmed and the nerve was allowed to fall back into its resting position. Again using the glass probe, the L-6 nerve was hooked from under the sacro-iliac rim, tightly ligated (6-0 silk), and placed back in position under the sacro-iliac rim. All damaged tissue was debrided, and all blood swabbed clear of the incision site. The muscle and tissue was gently rolled back into place to prevent any subdermal air pockets. The fascia over the muscle, and subsequently the skin was sutured shut (both using 3-0 Vicryl).

Post-operatively, the animals were placed in a recovery box with a heat lamp for observation. Following a successful surgery, the animals were fully ambulatory with no indications of spontaneous pain (licking, biting, or autonomy of the neuropathic foot, or vocalizations). If the animals presented initially with a foot drop
5 of the neuropathic paw, that disappeared within 1-2 days post-operatively. Any animals that presented with locomotor deficits or symptoms of spontaneous pain 2 days post-operatively, were promptly euthanised with an overdose of halothane.

B. Behavioral Testing

10

All behavioral tests were conducted between 08:00 and 14:00 daily, and occurred at 7, 12, 17, and 22 days following nerve injury. An exception to this was the cannulated animals, which were tested at 7 and 12 days after cannulation (14 and 19 days after nerve injury respectively). Following a recovery period of 7 days,
15 animals were moved from the vivarium, weighed, and acclimatized to the testing room for 40 minutes. After this initial period the animals were placed in the testing apparatus for 30-40 minutes or until exploratory behavior ceased.

B.1. Establishment and maintenance of neuropathic pain for the testing 20 periods

To determine that peripheral mononeuropathy resulted from ligation of L5 and L6 spinal nerves tests of static mechanical allodynia and thermal hyperalgesia were conducted.

25

B.2. Testing for Thermal Hyperalgesia

Surgically induced thermal hyperalgesia was assessed using a Paw Thermal Stimulator System. The animals were allowed two acclimatization periods prior to

the testing (30 minutes post removal from housing facility; 20 minutes in individual testing chambers). Rats were placed in pairs in a plexiglass box on top of the temperature maintained glass surface ($30 \pm 0.1^\circ\text{C}$) of the stimulator. After the initial acclimatization period, rats were tested for baseline withdrawal latencies (seconds) of both paws once every 20 minutes until three stable baselines were achieved. The animals were then returned to their cages for 30-40 minutes prior to drug administration and given ad-libitum access to food and water. Following drug administration, the animals were returned to the testing boxes for testing.

10 **B.3. Static Mechanical Allodynia**

Following weighing and testing room acclimatization, rats were placed in an elevated plexiglass container with a wire mesh bottom to allow access to the ventral surface of the hind paws. Rats were further acclimated to the testing chambers and tested for experimental baseline (two tests in one hour) using Semmes-Weinstein monofilaments (Stoelting Comp., Wood Dale IL.). The 50% withdrawal threshold was determined using the Dixons up-down method (Chaplan et al., 1994). Briefly, filaments were applied to the ventral surface of the paw starting with the 4.31 filament (2.04 gm) and the response noted. Following a positive response (paw withdrawal, with characteristic pain behavior), a lower filament was then applied. If no response was shown, the weight of the subsequent filament applied was increased. The 50% withdrawal threshold was then determined from the tabular value for the pattern of responses (k), the final monofilament value (X_f ; in log units), and interpolated using the formula:

$$25 \qquad 50\% \text{ g threshold} = (10^{[X_f + k\delta]}) / 10,000$$

where δ is the mean difference between stimuli. Following the determination of baseline values, the rats were returned to their original cages and left undisturbed with ad-libitum access to food and water for 30-40 minutes. After drug administration, the animals were again placed in the testing chambers and monitored for the appropriate

behavioral periods depending on the route of drug administration. The results of these tests are shown in Figure 1.

B.4. Data Analysis

5

For each of the experiments, raw data for response thresholds of both paws in each animal were recorded and entered into a spreadsheet (Microsoft Excel 5.0). The data was then normalized for each animal as Maximum Possible Effect (MPE) in terms of reversal of the neuropathic symptom being tested (MPE (anti-hyperalgesia or
10 antiallodynia)) or in terms of the analgesic effect of the contralateral paw (MPE (analgesia)). These values were calculated as follows:

$$\text{MPE (anti-neuropathic)} = (\text{PDR} - \text{IBR}) / (\text{CBR} - \text{IBR}),$$

wherein PDR is the post drug response of the ipsilateral paw, IBR is the ipsilateral paw baseline response, and CBR is the contralateral paw baseline response.

15 Similarly, the MPE (analgesia) was calculated as follows:

$$\text{MPE (analgesic)} = (\text{PDR} - \text{CBR}) / (\text{cutoff} - \text{CBR}),$$

wherein cutoff was 20 seconds from the thermal stimulus and 15 g for tactile allodynia, and PDR is the post drug withdrawal threshold of the contralateral paw.

Accordingly, the individual values reported or depicted are the $\text{MPE} \pm \text{SEM}$. The
20 time course of the drug effect is also depicted as the cumulative change in the MPE (ΔMPE) for each treatment and route of administration. This value was calculated as the sum of the individual MPE values through the time course (Σ_n), assuming an MPE value of 0 for the baseline values (not shown in figures). The data was statistically analyzed using a one way ANOVA for repeated measures with a pairwise comparison
25 (Dunnetts post hoc analysis). To compare the ΔMPE values of multiple treatment groups a one-way ANOVA with a pairwise comparison (Student-Neuman-Keuls post hoc analysis) was used. When comparing the means at individual time points or in the ΔMPE values, a student's T-test was used with a Student-Neuman-Keuls post hoc correction. For all tests, a P value less than .05 ($P < .05$) was considered significant.

These tests showed that spinal ligation resulted in neuropathic pain in rats with presenting symptoms of static mechanical allodynia and thermal hyperalgesia, and the condition was maintained throughout the testing periods (days 7, 12, 17, 22). As shown by the data summarized in Figure 1, the threshold values for thermal hyperalgesia were 7.98 ± 0.48 sec for the ipsilateral paw as compared to 10.35 ± 0.63 sec for the contralateral paw. Allodynia was manifested as 50% withdrawal threshold of 2.20 ± 0.28 g for the ipsilateral paw and 13.37 ± 0.37 g for the contralateral paw. Threshold values remained relatively stable with no statistically significant difference in baseline values for either the ipsilateral or the contralateral paw on each of the subsequent testing days.

B5. Drug Treatment

Tests were conducted to determine the antinociceptive effect of the tricyclic antidepressant amitriptyline in the ligated rat model. All drugs treatments were blinded. In both of the behavioral tests, amitriptyline (Sigma Chemical Co., St. Louis, MO) or saline was administered either systemically (by intraperitoneal (i.p.) injection), spinally (through an implanted intrathecal cannulas (i.t.)), or locally (by subcutaneous injection (s.c.) in the dorsal surface of the paw). Drugs were injected in volume of 5 ml/kg for systemic injections, and a total volume of 20 μ l (10 μ l drug + 10 μ l saline flush) for intrathecal injections. For local injections, the animals were briefly anaesthetized with halothane, and the solution was administered s.c. to the dorsal surface of the ipsilateral or contralateral paw in a volume of 50 μ l.

C1. Effect of systemic amitriptyline on nerve injury-induced thermal hyperalgesia

Systemic administration to nerve-injured rats of amitriptyline at doses of 1, 3 and 10 mg/kg of body weight by ip. injection produced an anti-hyperalgesic effect on the thermal threshold of the nerve-injured paw, as shown by the data summarized on Figures 2A-C. This effect was maximal at 10 mg/kg; reaching an almost complete reversal of thermal hyperalgesia ($MPE = 0.94 \pm -0.17$) 60-80 minutes after injection (Figure 2A). A slight but not statistically significant elevation in threshold was observed in the contralateral paw for the 10 mg/kg dose (Figure 2B). Comparison of the ΔMPE values for the ipsilateral paw (Figure 2C), shows a statistically significant antinociceptive effect of 10 mg/kg for amitriptyline as compared to saline controls, as well as for 1 mg/kg and 3 mg/kg doses. While both 1 mg/kg and 3 mg/kg doses produced results significantly different from those of the saline controls, they were not significantly different from each other. A slight sedative effect of the 10 mg/kg dose was observed in a few of the rats, but none was observed at other doses. This effect did not impact the ability of the animals to respond to the stimulus, as the vigor of the response at threshold was similar to that observed during baseline determinations.

C2. Intrathecal Cannulations

In animals used to study the effects of spinal administration of amitriptyline, intrathecal cannulas were implanted 7 days following nerve injury. For the implantation of intrathecal cannulas for the spinal delivery of drugs, rats were anaesthetized with halothane and mounted in the earbars of a stereotaxic apparatus. The cisternal membrane at the base of the skull was exposed, a small incision made, and a cannula (7.5 cm of PE-10 tubing) inserted into the subarachnoid space. The cannula was filled with saline and the external end was occluded with a small wire plug. Rats were injected with Penlong XL^R (0.15ml i.m.) and lactated Ringers

solution (10 ml s.c.) while still anaesthetized. On the day of testing, rats were accommodated in a plastic container for the intrathecal injection (in a 10 μ l volume followed by 10 μ l saline to flush the cannula).

5 **C3. Effect of spinal amitriptyline on nerve-injury induced thermal hyperalgesia**

The intrathecal injection of amitriptyline (60 μ g) resulted in a partial reversal of the thermal hyperalgesia of the ipsilateral paw in nerve injured rats that was evident 10 20 minutes after injection and lasted for the 60 minute testing period, as shown by the data in Figures 3A-C. Because a 30 μ g dose was ineffective in the allodynia test, and 90 μ g was highly sedative, these doses were not used for testing of the thermal hyperalgesic effect. While values for the individual time point are not significantly different on their own (Figure 3A), the Δ MPE values for saline and amitriptyline 15 (60 μ g) were significantly different (Figure 3C). No significant difference was observed, in the contralateral paw withdrawal thresholds, at either single time points or in comparison of the Δ MPE values (Figures 3B and 3C).

20 **C4. Effect of local injection of amitriptyline on nerve injury-induced thermal hyperalgesia**

When injected locally into the neuropathic paw, amitriptyline at a dose of 100 nmol, which produces a local antinociceptive effect in the rat formalin test as described in Example 2 herein (Figure 8), had an immediate statistically significant 25 antinociceptive effect, almost completely reversing the thermal hyperalgesia in nerve-injured rats (Figure 4A). The MPE values remained significantly different from those of saline for the first 120 minutes. A contralateral local injection of amitriptyline (100 nmol) failed to show any significant effect on the withdrawal latency of the ipsilateral paw (Figure. 4A), a result that indicates a lack of systemic effect. No local

effect of the injected paw was observed either for the control rats injected with saline (ipsilateral paw) or for animals injected with amitriptyline (100 nmol) in the contralateral paw.

5 There was also no significant alteration in the withdrawal thresholds of naive animals after local injection of amitriptyline 100 nmol (n=4; data not shown). In addition, neither the ipsilateral nor contralateral local injections showed any significant effect on the contralateral paw withdrawal latency (Figure 4B). Comparison of the Δ MPE values revealed a statistically significant difference
10 resulting from locally injected amitriptyline on thermal hyperalgesia as compared to both the saline control and the contralateral injection control (Figure 4C).

C5. Effect of systemic amitriptyline on nerve injury-induced mechanical allodynia

15 Amitriptyline was systemically administered to nerve injured rats at doses of 1.5, 5.0, and 10.0 mg/kg of body weight. No significant effect on nerve injury induced allodynia in the ipsilateral paw was observed at any of the doses (Figure 5A). Since 10 mg/kg was slightly sedative in some rats, a higher dose was not used. In the
20 contralateral paw, all doses of amitriptyline caused a hyperaesthetic response resulting in a reduced MPE (analgesia) value ranging from -0.4 to -0.8 (Figure 5B). The Δ MPE values for each dose differed significantly from saline (Figure 5C). Qualitatively, the hyperaesthetic response was observed as a brisk withdrawal to the filament with no overt signs of typical neuropathic pain responses such as biting,
25 licking, vocalization or guarding of the paw. Although the values of the response threshold in the non-injured paws were very similar to those in the injured paws, the responses were not categorized as 'neuropathic pain responses'. Threshold testing 24 hours after drug administration showed that both the ipsilateral and contralateral baselines returned to predrug values (data not shown).

C6. Effect of spinal amitriptyline on nerve injury-induced mechanical allodynia

5 Amitriptyline was also administered spinally to test its antinociceptive effect against nerve injury-induced mechanical allodynia in the ipsilateral paw. At doses of 30, 60, and 90 μg administered spinally, amitriptyline failed to have a significant effect on allodynia in the ipsilateral paw (Figures 5A and 6C). However a comparison of the ΔMPE for the three groups revealed the 60 μg group to be significantly
10 different from the 30 μg and 90 μg groups, as well as from the saline group (Figure. 6C). Spinally administered amitriptyline (60 μg) caused a hyperaesthetic response in the contralateral paw (Figures 6B and 6C) that was similar in magnitude to that observed following systemic administration. While 90 μg dose ($n=3$) appeared not to facilitate the hyperaesthetic response, this result was probably due largely to the
15 pronounced sedative effect of the drug at this dose, which interfered with accurate determination of the withdrawal threshold.

C7. Effect of local amitriptyline on nerve-injury-induced mechanical allodynia

20 Amitriptyline was also injected locally to test the local effect on nerve-injury induced mechanical allodynia. Local injection of amitriptyline (100 nmol) failed to show any significant effect on the withdrawal response of the nerve injured paw to tactile stimulation (Figures 7A and 7C). Local injection of amitriptyline into the
25 nerve injured paw did, however, cause a reduction in the response threshold of the contralateral paw. This effect also occurred after direct local injection of amitriptyline (100 nmol) into the non-injured contralateral paw (Figure 7B). The response to locally administered drug in the contralateral paw was observed to be significantly different from the response to saline, but not different from the response following

injection into the non-injured paw at individual time points (Figure 7B) and in comparison of the Δ MPEs (Figure 7C).

EXAMPLE 2

5

Experiments were performed using male Sprague-Dawley rats (120-160 g; group mean 142 ± 4 g) (Charles River, Quebec, Canada). Rats were housed in pairs and maintained on a 12:12 h light/dark cycle at $22 \pm 1^\circ\text{C}$. Food and water were freely available.

10

The test drugs were amitriptyline, caffeine, 5'-amino-5'-deoxyadenosine (NH_2dAD), phentolamine HCl, dimethylsulfoxide (DMSO) (Sigma Chemical Co., St. Louis, MO), 8-cyclopentyl-1,3-dimethylxanthine (CPT), N^6 -benzyl-N-ethylcarboxamide adenosine (N^6 -benzyl-NECA), MK-801 hydrogen maleate, naloxone HCl (Research Biochemicals Inc., Natick, NJ), 2'-Deoxycoformycin (Parke Davis Pharmaceutical Research Division, Ann Arbor, MI). Formalin (37% formaldehyde) (British Drug Houses, Toronto, Ontario) was the control irritant. All drugs except for CPT were dissolved in saline and diluted to the appropriate dose with formalin. CPT was dissolved in a final concentration of 15% DMSO, and appropriate control groups were included.

20

A. The Formalin Test

On the day of testing, rats were placed in a 28x28x28 cm plexiglass observation chamber for an initial 20 min to allow familiarization with surroundings. The indicated doses of formalin (0.5-2.5%) and formalin/drug combinations were injected s.c. in a volume of $50\mu\text{l}$ into the dorsal hindpaw of the rat. To establish whether inhibition of pain response was due to a systemic rather than a local effect, the test drug was administered into the contralateral paw. For injections into the

25

contralateral paw, the drug was injected immediately before the formalin in the other paw. Following injections, rats were returned to the observation chamber, and flinching behaviors (lifting, shaking or rippling of the haunch) were monitored over two time periods, phase 1 (0-12 minutes following administration) and phase 2 (16-60 minutes following administration). Biting/licking time was also monitored. Flinches, which were rapid events, were recorded as discrete episodes, while the time spent biting or licking was recorded cumulatively. Two rats in adjacent chambers were observed at a time, with observations taking place in alternating 2 min bits. Recorded episodes and times were not corrected, with the result that values obtained represent about half of the total behaviors expressed.

Data was analyzed using analysis of variance followed by the Student-Neuman-Keuls test for multiple groups, or the Student t-test for two groups.

B. Local Antinociceptive Action of Amitriptyline

To test the antinociceptive effect of locally administered antidepressant, amitriptyline was coadministered at doses of 10, 30, and 100 nmol with 2.5% formalin. A dose-dependent reduction in the expression of flinching behaviors both in phase 1 and phase 2 following injection was observed. The time course for these effects is presented in Figure 8A, and the cumulative scores for the two phases are presented in Figures 9A and 9B. A similar dose-related inhibition in biting/licking behaviors was obtained (Figures 8B and 9C); however, there was no distinct phase 1 effect observed in biting/licking behaviors (Figure 8B). The inhibitory effect of amitriptyline on both parameters was not observed following administration of amitriptyline into the contralateral hindpaw (Figures 9A-B). The absence of inhibitory effect on the contralateral hindpaw indicates that the antinociceptive action of amitriptyline is a local effect.

Inhibition of pain was seen most clearly at higher concentrations of formalin. For example at 0.75% formalin, the effect was seen only with flinching behaviors, while at 0.5%, it was not seen at all (Figure 3).

5 C. Involvement of Adenosine in the Local Antinociceptive Action of Amitriptyline

To examine involvement of adenosine in the antinociceptive action of amitriptyline, caffeine, a non-selective adenosine antagonist, was coadministered at
10 doses of 500 and 1500 nmol with amitriptyline. A significant reversal of the antinociceptive action of amitriptyline was observed in phase 2 flinches and total biting/licking time, but phase 1 flinches were not affected (Figures 11A-C). Although there was only a partial reversal of amitriptyline action at 2.5% formalin caused by caffeine, caffeine clearly reversed the local effects of adenosine. Therefore, it was
15 speculated that the expression of adenosine may depend on the intensity of the pain stimulant. To test this hypothesis, some experiments were undertaken using a lower concentration of formalin. Amitriptyline at 100 nmol concentration reduced phase 2 flinches to formalin 1.5% to a lesser extent than in the 2.5% formalin test, but caffeine produced a complete reversal of the antinociceptive action of amitriptyline
20 (Figure 13A). However, at a higher dose of amitriptyline (300 nmol), which produced a comparable extent of inhibition as did 100 nmol with 2.5% formalin, the degree of reversal by caffeine was still only partial (Figure 13A).

CPT, a selective adenosine A₁ receptor antagonist, coadministered at a dose of
25 150nmol with 100 nmol amitriptyline produced a similar degree of reversal of the action of amitriptyline as did caffeine in the 1.5% formalin test (Figure 13B). At 1.5% formalin, the effect of amitriptyline on phase 1 flinches was not reversed in any experiment, while the effect on biting/licking was reversed only with the caffeine at 100 nmol amitriptyline (data not shown). Caffeine and CPT did not produce an

intrinsic effect in the 1.5% formalin test (Figure 13A-B), but caffeine produced a slight reduction in flinches at 2.5% formalin concentration as can be seen by comparing the data of (Figures 11A-C and Figures 12A-B).

- 5 Coadministration of NH_2dAD , an inhibitor of adenosine kinase, at a dose of 100 nmol augmented the action of a partially effective dose of amitriptyline on phase 2 flinches in a 2.5% formalin test (Figure 14), but was without effect on phase 1 flinches and biting/licking time (data not shown). This dose of NH_2dAD is without intrinsic effect on formalin behaviors at this concentration of formalin.

10

D. Involvement of Other Mechanisms in the Local Antinociceptive Action of Amitriptyline

Amitriptyline is known to exert a number of pharmacological effects.

- 15 Experiments were conducted to test for potential involvement of interactions with excitatory amino acid receptors, biogenic amines (noradrenaline, 5-hydroxytryptamine or 5-HT, histamine) and endogenous opioids by determining effects of respective receptor antagonists. The coadministration of the non-competitive NMDA receptor antagonist, MK-801 at doses of 20 and 100 nmol with formalin produced a dose-
20 dependent reduction in phase 2 flinching behaviors and biting/licking time (Figures 15A-B), with the latter behavior exhibiting a greater sensitivity to inhibition. No significant effects were seen on phase 1 flinching behaviors (data not shown). This action of MK-801 is not locally mediated, as shown by the identical effect seen following administration of the MK-801 into the contralateral hindpaw
25 (Figures 15A-B). These results indicate that MK-801 acts by a systemic route. Injection of 100 nmol (but not 20 nmol) into both the ipsilateral and contralateral hindpaws also produces a behavioral hyperactivity during the observation interval. Lower doses of MK-801 (0.03 and 0.3 nmol) coadministered with formalin had no effect on any parameter (n=4 each, data not shown).

Coadministration of phentolamine (PTA), a non-selective-adrenoreceptor antagonist, had no significant effect on the action of amitriptyline at 100 nmol, but coadministration of 500 nmol augmented its action (Figure 16A). This potentiation is due to an additive action of an intrinsic effect of phentolamine with that of amitriptyline, as phentolamine alone at this dose inhibited the action of formalin (Figure 16A). An identical pattern of interactions was seen with the phase 2 biting/licking time, but no significant effects were seen with phentolamine on phase 1 flinching (data not shown).

To examine a potential local antihistamine or anti-serotonergic action for amitriptyline, tests were designed to determine whether amitriptyline could alter the flinching behaviors produced by N^6 -benzyl NECA, an adenosine A_3 receptor agonist which produces intrinsic nociceptive behaviors by releasing histamine and 5-HT from mast cells. In these tests, 100 nmol amitriptyline did not alter the number of flinches generated by 5 nmol N^6 -benzyl-NECA (25.5 ± 6.3 flinches in the absence, and 27.8 ± 4.3 flinches in the presence of amitriptyline, $n=6$ per group). Naloxone, an opioid receptor antagonist, did not alter the local effect of amitriptyline against phase 2 flinching (Figure 16B), phase 1 flinching, or biting/licking time (data not shown).

EXAMPLE 3

The drugs tested were desipramine, fluoxetine, caffeine and propranolol (Sigma Chemical Co., St. Louis, MO), ketanserin and tropisetron (Research Biochemicals Inc., Natick, NJ), and GR113808A (Glaxo Research Group, Greenford, Middlesex, U.K.). Formalin (37% formaldehyde) (British Drug Houses, Toronto, Ontario) was the control irritant. All drugs were prepared in saline and diluted to the indicated dose with formalin when coinjected.

Rats were maintained and surgically altered by spinal nerve ligation of the 5th and 6th spinal nerves as described in Example 1. Preliminary tests showed that the spinal ligation surgery generates a thermal hyperalgesia of 2-3 sec in the paw corresponding to the ligation (baseline on non-ligated side 10-12 sec, and 710 on ligated side), and this effect was maintained for at least 22 days following surgery.

A. The Formalin Test

The formalin test was applied as described in Example 2 above. Following injections, rats were returned to the observation chamber and scored for flinching behaviors (lifting, shaking or rippling of the haunch) and biting/licking time. Two rats in adjacent chambers were observed at a time, with observations occurring in alternate 2 min bits. Recorded episodes were not corrected for this, so values represent about half of the total behaviors expressed.

B. Local Antinociceptive Action of Desipramine and Fluoxetine

When independently injected with formalin 2.5% into the dorsal hindpaw of a non-surgically altered rat, desipramine (10-300 nmol) and fluoxetine (30-300 nmol) both produced a dose-dependent antinociceptive action against phase 2 flinching behaviors (Figure 17B) and phase 2 biting/licking behaviors (Figure 17C). Phase 1 flinching behaviors exhibited much more variability, and were only reduced significantly at the highest doses (Figure 17A). There was no phase 1 expression of biting/licking behaviors (Figure 18C). Desipramine action suppressed both pain response behaviors (illustrated in Figures 18A-B) in a uniform manner throughout the entire time course observed. A similar pattern of suppression of behaviors was observed with fluoxetine (data not shown). However, when the doses that were most efficacious in the ipsilateral paw were injected into the contralateral hindpaw, the antinociceptive action resulting from coadministration of desipramine or fluoxetine

with formalin was not observed. (Figures 17A-C, 18B and 18C). These results show that the antinociceptive effect of the locally injected antidepressant was local only.

C. Coadministration of Desipramine or Fluoxetine and Caffeine

5

To test the effect of caffeine on the antinociceptive effect of the antidepressant drugs, a formalin concentration of 1.5% was used as this concentration revealed the involvement of adenosine in the action of amitriptyline most clearly. (See Example 2 above). Coadministration of caffeine (1500 nmol) with desipramine or fluoxetine had
10 no significant effect on the antinociceptive actions of the antidepressants in any aspect of the formalin test (Figures 20A-C).

D. Local Administration of Desipramine and Fluoxetine in thermal hyperalgesia test

15

Nerve damaged rats (SNL) were anaesthetized briefly for drug injections (in 50 µl) into the dorsal part of the hindpaw, either on the side ipsilateral to the surgery (ligated side) to test for local effects, or the contralateral side to test for systemically mediated effects. Thermal thresholds were determined with the investigator unaware
20 of the particular treatment.

The local administration of desipramine (100 nmol), but not fluoxetine (100 nmol), into the hindpaw corresponding to the operated side produced a local anti-hyperalgesic effect when reaction thresholds were determined using a thermal
25 paw stimulator (Figures 19A and 4B). This effect was not observed following injection into the contralateral paw corresponding to the non-operated side (Figure 19B) These results show that the antinociceptive effects observed in the ipsilateral side were local. The local injection of desipramine or fluoxetine, 100 nmol each, into a normal rat paw (control) had no effect on thermal thresholds (n=4 each,

data not shown). Coadministration of caffeine (1500 nmol), which inhibits the local action of amitriptyline on thermal hyperalgesia following spinal nerve ligation (Example 1), also blocked the antihyperalgesic action of desipramine in the spinal nerve ligation model (Figures 21A-C).

5

E. Effects of Desipramine, Fluoxetine and Amitriptyline on Paw Volume

Paw volume was determined by plethysmometry using a commercially available device (Ugo, Basile). The hindpaw was inserted into the solution to the junction of the hairy and non hairy skin, and volume was determined by displacement. Triplicate baseline determinations were determined prior to the formalin injection, and again immediately following the behavioral observation interval, which lasted 60 min. For time course determinations, volumes were determined at 30 min intervals as indicated up to 3 hours in separate groups of rats.

15

The anti-inflammatory effect of desipramine and fluoxetine was tested by the independent coadministration of desipramine and fluoxetine (doses of 100 or 300 nmol) with formalin. No significant effect on the increase in paw volume induced by 1.5% formalin alone was observed by the antidepressants, although there was a tendency towards an increased paw volume resulting from fluoxetine (Figure 22A) at 2.5% formalin. When administered into the contralateral hindpaw, fluoxetine also significantly increased the paw volume in the absence of an inflammatory stimulus (Figure 22B). The time course of the actions of desipramine and fluoxetine was also determined. In these tests desipramine produced a limited effect on paw volume (Figure 23A), but fluoxetine produced a clear dose-related increase in paw volume that was maintained for the entire 3 hr observation interval (Figure 23B). The increase in paw volume was not accompanied by a reddening of the paw, or by spontaneous behaviors indicative of discomfort (e.g. lifting, licking, shaking). Amitriptyline also produced a limited increase on paw volume (Figure 23C), and did

25

not significantly alter the degree of paw swelling produced by formalin 1.5% (data not shown).

**F. Effects of Receptor Subtypes in Nociceptive Action of Desipramine and
5 Fluoxetine**

To test for involvement of 5-HT receptor subtypes in the action of fluoxetine (100 nmol), the effects of propranolol (5-HT₁ receptor antagonist) (300 nmol), ketanserin (5-HT₂ receptor antagonist) (500 nmol), tropisetron (5-HT_{3/4} receptor
10 antagonist) (300 nmol), and GR113808A (5-HT₄ receptor antagonist) (500 nmol) on paw volume were determined at doses known to block 5-HT actions when injected into the hindpaw (G.J. Doak and J. Sawynok (1997) *Neuroscience* 80:939-949). The results of these tests show that the increase in paw volume induced by fluoxetine was blocked by ketanserin but not by any of the other antagonists, indicating an
15 involvement of 5-HT₂ receptor subtypes in this action (Figures 24A-B).

EXAMPLE 4

**A. The Role of 5-HT₂ Receptor Antagonism in Peripheral Analgesia by
20 Amitriptyline**

Studies were conducted to resolve an apparent inconsistency in the art regarding the mechanism of analgesia of amitriptyline and other tricyclic, second generation, or third generation antidepressants of the invention. H. Hall et al. first
25 suggested that a number of antidepressants can block 5-HT₂ receptors ((1984) *Acta Pharmacol Toxicol*, 54:379-384). More recently F.V. Abbott et al. demonstrated that peripheral 5-HT₂ receptor blockers can block the expression of formalin-evoked behaviors using the formalin test as described above, except that 1% formalin was used and proposed based on the results of this low dose study that 5-HT₂ receptor
30 antagonists might make useful analgesics ((1996) *Neuropharmacology* 35:99-110).

However, when Doak and Sawynok, supra (1997) conducted a detailed investigation into the 5-HT receptor subtypes involved in formalin-evoked behaviors using a 2.5% formalin test, evidence was obtained for the involvement of 5-HT₁, 5-HT₃ and 5-HT₄ but not 5-HT₂ receptors in such behaviors. Figure 25 above summarizes the results of the Doak and Sawynok studies of 1997.

A number of drug effects are known to be intensity-dependent (in their expression) in the formalin test (e.g., Sawynok J., Reid AR (1996) *Eur J Pharmacol*, 298:105-111). In view of the apparent contradiction between the results obtained in the Abbott study in which 1% formalin was used and in the Doak and Sawynok study, in which 2.5% formalin was used, studies were undertaken to re-evaluate the performance of ketanserin, the selective 5-HT₂ receptor antagonist used in both studies.

As the Abbott study used 1% formalin rather than the 2.5% formalin used in the Doak and Sawynok study, the lower dosage of formalin was used to re-evaluate the performance of ketanserin using individual groups of male Sprague Dawley rats (n=6 per group) to conduct the 1% formalin test as in Example 2 A. above. The results of the re-evaluation shown in Figure 26 above indicate ketanserin at doses of 50 nmol and 500 nmol (a dose equal to or greater than the 9.5-50 µg or 0.92 - 92 nmol used in the Abbott study) fails to block formalin-evoked flinches in Sprague Dawley rats even at 1% formalin. By contrast, co-administration of 3 nmol and 30 nmol doses of ketanserin were used to blockade change in paw volume induced by local injection of 5-HT into the hindpaw of Sprague Dawley rats. Drugs were injected in a volume of 50 µl and paw edema was measured by plethysmometry (volume displacement). The results of these studies shown in Figures 27 A and B clearly indicate that the dosage of ketanserin that failed to prevent 1% formalin-induced flinching behavior was sufficient to block the paw edema produced by local injection of 5-HT.

The only other difference between the Abbott study and the Doak and Sawynok study was the species of rat used, Long Evans rats in the former and Sprague Dawley rats in the latter. It may be that there is a differential expression of 5-HT receptors on sensory afferent nerve terminals in the two species. However, it should be noted that
5 in the absence of any effect of the 5-HT₂ antagonist in the test in species and conditions where a profound analgesia is seen with amitriptyline (2.5% formalin, Sprague Dawley rats), it is not shown that 5-HT₂ antagonism accounts for the analgesia demonstrated by amitriptyline and other tricyclic, second generation, or third generation antidepressants observed in the studies described herein.

10

B. The Role of Blockade of Cholinergic Muscarinic and Nicotinic Receptors in Peripheral Analgesia by Amitriptyline

Amitriptyline is known to bind to central muscarinic cholinergic receptors
15 (Hall H, Ogren SO. (1981) *Eur J. Pharmacol*, 70:393-407; Choi A, Mitchelson F. (1994) *Pharmacology*, 48:293-300), and its affinity for such receptors has been implicated in certain side effects of amitriptyline observed clinically (Baldessarini R.J. (1990) *The Pharmacological Basis of Therapeutics*, 383-435). Amitriptyline also blocks nicotinic receptor-mediated responses (Schofield G.G. et al. (1981) *Proc Natl*
20 *Acad Sci USA*, 78:5420-5244; Fryer J.D., Lukas R.J. (1999) *J Neurochem*, 72:1117-1124), although in this case the block seems to result from an inhibition of the ion channel itself rather than a competitive inhibition at the receptor binding site. Acetylcholine can stimulate sensory nerve endings to produce pain (Steen K.H., Reeh P.W. (1993) *J Neurophysiol*, 70:397-405; Bauer M.B. et al. (1994) *Neuroscience*,
25 62:351-359), and both nicotinic and muscarinic receptors are present in dorsal root ganglion neurons and on sensory nerve endings (Ninkovic M., Hunt S.P. (1983) *Brain Research*, 272:57-69; Wamsley J.K. et al. 1981) *Brain Research*, 217:155-161). In view of these findings, studies were conducted to determine whether established

antagonists for muscarinic and nicotinic cholinergic receptors can mimic the action of amitriptyline.

In these studies, groups of male Sprague Dawley rats (n=5-8 per group) received doses ranging from 10 to 1000 nmol of atropine (muscarinic receptor antagonist), mecamylamine (ganglionic nicotinic receptor antagonist) and d-tubocurarine (neuromuscular nicotinic receptor antagonist) coadministered with formalin in the formalin 2.5% test. Figures 28A-C show the effects of the drugs on phase 1 (Figure 28A) and phase 2 flinching (Figure 28B) and phase 2 biting/licking behaviors (Figure 28C), which were determined as described above.

In these tests, atropine produced a reduction in flinching behaviors only at the highest dose tested (1000 nmol). This effect is a local peripheral action, as it was not observed following administration of atropine into the contralateral hindpaw. Atropine also produced a dose-dependent reduction in biting/licking behaviors, and this appeared to be a composite of peripheral and central actions as administration of atropine into the contralateral hindpaw also produced a reduction in this behavior at the highest dose (Figure 28C). Mecamylamine had no significant effect on either behavior at the highest dose. d-Tubocurarine had no effect on formalin-evoked behaviors at 10 nmol, and at 100 nmol, d-tubocurarine was lethal, probably due to systemic activity of the drug on respiratory muscles causing respiratory arrest.

These results indicate that established cholinergic receptor antagonists are only weakly active (atropine) or are inactive (mecamylamine, d-tubocurarine) at inhibiting formalin-evoked behaviors when administered peripherally. In view of these results, it can be concluded that the actions of amitriptyline on formalin behaviors are due to blockade of peripheral cholinergic receptors on sensory nerve terminals.

C. Conclusions

The present results provide no evidence to support an involvement of blockade of peripheral 5-HT₂, muscarinic cholinergic or nicotinic cholinergic receptors in the analgesic actions of amitriptyline in the rat formalin test. To date, despite considering the possibility of a potential involvement of antihistaminic, antiadrenergic, antiserotonergic and anticholinergic actions in the action of amitriptyline, the only mechanism clearly implicated in the action of amitriptyline is some form of interaction with adenosine receptors as peripheral analgesia is partially blocked by the adenosine receptor antagonist caffeine.

It will be apparent to those skilled in the art that various changes may be made in the invention without departing from the spirit and scope thereof, and therefore, the invention encompasses embodiments in addition to those specifically disclosed in the specification, but only as indicated in the appended claims.

That which is claimed is:

1. A method for producing local analgesia in a subject having a site of local discomfort, said method comprising locally administering an effective amount of a tricyclic, second generation or third generation antidepressant to the site.

2. The method according to claim 1 wherein the antidepressant is a tricyclic antidepressant.

3. The method according to claim 1 wherein the tricyclic antidepressant has a structure:



wherein Z is a 7-membered ring, optionally containing 1 or 2 biocompatible
5 heteroatoms, or an 8-membered bicyclic ring,

Ar₁ and Ar₂ are optionally substituted aromatic rings fused to Z, and

R is an alkylamino or arylamino substituent, optionally an N-oxide derivative thereof.

4. The method according to claim 3 wherein the heteroatom is oxygen or nitrogen.

5. The method according to claim 3 wherein Ar₁ and Ar₂ are independently optionally substituted with a biocompatible halogen.

6. The method according to claim 3 wherein the alkylamino or arylamino substituent is an N-oxide derivative thereof.

7. The method according to claim 3 wherein the alkylamino substituent comprises from 4 to 5 carbons atoms.

8. The method according to claim 7 wherein the alkylamino is a tertiary or secondary amino group.

9. The method according to claim 3 wherein R is selected from the group consisting of


- 5
- $(\text{CH}_2)_3\text{N}(\text{CH}_3)_2$,
 - $(\text{CH}_2)_3\text{NHCH}_3$,
 - $(\text{CH}_2)_3\text{N}(\text{CH}_3)_2$,
 - $\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{N}(\text{CH}_3)_2$,
 - = $\text{CH}(\text{CH}_2)_2\text{N}(\text{CH}_3)_2$,
 - $(\text{CH}_2)_3\text{NHCH}_3$, and
 - = $\text{CH}(\text{CH}_2)_2\text{NHCH}_3$.

10. The method according to claim 1 wherein the tricyclic antidepressant is selected from the group consisting of clomipramine, imipramine, amitriptyline, doxepin, desipramine, nortriptyline, amoxapine, maprotiline, trimipramine, and suitable combinations of any two or more thereof.

11. The method according to claim 1 wherein the tricyclic antidepressant is amitriptyline or desipramine.

12. The method according to claim 1 wherein the second generation or third generation antidepressant is selected from the group consisting of mirtazapine, venlafaxine, trazodone, bupropion, fefazodone, and suitable combinations of any two or more thereof.

13. The method according to claim 12 wherein the second generation or third generation has the structure



wherein Ar_3 is an N-containing heterocyclic ring,

- 5 Y is either an aryl group fused to the heterocyclic ring or one or two substituents selected from alkyl, arylalkyl, arylalkyloxy, aryl and heteroaryl substituents comprising a total of about 4 to 8 carbons attached to Ar_3 , and attached to Q at a second nitrogen atom of Ar_4 .

14. The method according to claim 13 wherein the X is an alkyl group containing 3 carbons.

15. The method according to claim 13 wherein Ar_3 is a 1,2,4-triazole, Q is an heteroarylalkyl substituent containing 6 to 8 carbon atoms, and is substituted at the 4 position with Q.

16. The method according to claim 13 wherein the alkyl substituent contains 3 carbon atoms.


17. The method according to claim 13 wherein the benzene ring is substituted with a halogen selected from the group consisting of chlorine, bromine, and fluorine.

18. The method according to claim 1 wherein the tricyclic-antidepressant is desipramine.

19. The method according to claim 1 wherein the tricyclic-antidepressant is amitriptyline.

20. The method according to claim 1 wherein the tricyclic-antidepressant is administered topically.
21. The method according to claim 1 wherein the antidepressant is administered as a formulation selected from the group consisting of a cream, a lotion, a gel, an ointment, a spray, a polymer stabilized crystal, a powder, and an aerosol.
22. The method according to claim 17 wherein the antidepressant is administered by a method selected from injection, supersonic powder injection, and transdermal electroporation.
23. The method according to claim 1 wherein the antidepressant is administered encapsulated in a delivery system selected from the group consisting of a microsphere, a polymer stabilized crystal, and a liposome.
24. The method according to claim 1 wherein the analgesia suppresses pain caused by inflammation.
25. The method according to claim 1 wherein the analgesia suppresses neuropathic pain.
26. A composition for local administration comprising a tricyclic or heterocyclic antidepressant other than doxepine, and a vehicle suitable for topical administration.
27. The composition according to claim 26 wherein the antidepressant is a tricyclic antidepressant.

28. The composition according to claim 26 wherein the tricyclic antidepressant has a structure:



wherein Z is a 7-membered ring, optionally containing 1 or 2 biocompatible
5 heteroatoms, or an 8-membered bicyclic ring,

Ar₁ and Ar₂ are optionally substituted aromatic rings fused to Z, and

R is an alkyl amino or aryl amino substituent, and optionally an N-oxide derivative thereof.

29. The composition according to claim 28 wherein the heteroatom is oxygen or nitrogen.

30. The composition according to claim 28 wherein Ar₁ and Ar₂ are independently optionally substituted with a biocompatible halogen.







31. The composition according to claim 28 wherein the alkylamino or arylamino substituent is an N-oxide derivative thereof.

32. The composition according to claim 28 wherein the alkylamino substituent comprises from 4 to 5 carbons atoms.

33. The composition according to claim 32 wherein the alkylamino is a tertiary or secondary amino group.

34. The composition according to claim 28 wherein R is selected from the group consisting of


5



35. The composition according to claim 26 wherein the tricyclic antidepressant is selected from the group consisting of clomipramine, imipramine, amitriptyline, desipramine, nortriptyline, amoxapine, maprotiline, trimipramine, and suitable combinations of any two or more thereof.

36. The composition according to claim 26 wherein the tricyclic antidepressant is amitriptyline or desipramine.

37. The composition according to claim 26 wherein the second or third generation antidepressant has a structure:



wherein Ar_3 is a substituted N-containing heterocyclic ring,

5 Y is either an aryl group fused to the heterocyclic ring, or one or two substituents selected from the group consisting of alkyl, alkyloxy, arylalkyl, arylalkyloxy, aryl, heteroaryl substituents, and combinations thereof comprising a total of about 4 to 8 carbons attached to Ar_3 ,

X is an alkyl group comprising 2 to 5 carbon atoms linking Ar_3 and Ar_4 ,

10 Ar_4 is a piperazine attached to X by a first nitrogen atom of Ar_4 , and

Q is a benzene ring optionally substituted with a biocompatible halogen and attached to Ar_4 at a second nitrogen atom of Ar_4 .

38. The composition according to claim 37 wherein the X is an alkyl group containing 3 carbons.

39. The composition according to claim 37 wherein Ar_3 is a 1,2,4-triazole substituted at the 4 position with the arylalkyloxy substituent containing 6 to 8 carbon atoms.

40. The composition according to claim 39 wherein the heteroarylalkyl substituent contains an oxygen atom.

41. The composition according to claim 37 wherein the benzene ring is substituted with a halogen selected from the group consisting of chlorine, bromine, and fluorine.

42. The composition of claim 23 further comprising an inert carrier.

43. The composition of claim 42 wherein the inert carrier is selected from the group consisting of water, isopropyl alcohol, gaseous fluorocarbons, ethyl alcohol, polyvinyl pyrrolidone, propylene glycol, a fragrance, a gel-producing material, stearyl alcohol, stearic acid, spermaceti, sorbitan monooleate, methylcellulose, and suitable combinations thereof.
44. The composition according to claim 26 wherein the composition further comprises a penetration enhancing agent.
45. The composition according to claim 26 wherein the tricyclic-antidepressant is selected from the group consisting of desipramine, amitriptyline and a suitable combination thereof.
46. The composition according to claim 26 wherein the composition further comprises a penetration enhancing agent.
47. The composition according to claim 26 wherein the tricyclic-antidepressant is desipramine.
48. The composition according to claim 26 wherein the tricyclic-antidepressant is amitriptyline.
49. The composition according to claim 26 in a formulation selected from the group consisting of a cream, a lotion, a gel, an ointment, a spray, a powder, a polymer stabilized crystal, and an aerosol.
50. The composition of claim 26 further comprising a neutralizing agent.

51. The composition of claim 26 wherein the composition is formulated for local injection.

52. The composition according to claim 26 wherein the antidepressant is encapsulated in a slow release delivery vehicle.

53. The composition according to claim 52 wherein the delivery vehicle is selected from a liposome, a microcapsule, a polymer stabilized crystal.

1/29







FIGURE 1

2/29



3/29




FIGURE 3A




FIGURE 3B




FIGURE 3C

4/29




FIGURE 4A




FIGURE 4B




FIGURE 4C

5/29

A. Ipsilateral Paw




FIGURE 5A

B. Contralateral Paw




FIGURE 5B





C. Cumulative MPE (Σ MPE)

FIGURE 5C

6/29



7/29




FIGURE 7A




FIGURE 7B




FIGURE 7C

FIGURE 8A





FIGURE 8B



9/29

FIGURE 9C




FIGURE 9B







FIGURE 9A



10/29

FIGURE 10A



11/29




FIGURE 11A

FIGURE 11B

FIGURE 11C






FIGURE 12B


FIGURE 12A

13/29



14/29

FIGURE 14



15/29




FIGURE 15A

FIGURE 15B

16/29

FIGURE 16A





FIGURE 16B



17/29

FIGURE 17A





FIGURE 17B









FIGURE 17C




18/29

A. IPSILATERAL**FIGURE 18A****B. CONTRALATERAL****FIGURE 18B**

19/29


C. IPSILATERAL

FIGURE 18C






D. CONTRALATERAL

FIGURE 18D



20/29



21/29




FIG. 20A







FIG. 20B

22/29









FIGURE 22A

FIGURE 22B

24/29



25/29

FIGURE 24B




FIGURE 24A





FIGURE 25



27/29

FIGURE 26




FIGURE 27A





FIGURE 27B



29/29

FIGURE 28A




FIGURE 28B





FIGURE 28C



INTERNATIONAL SEARCH REPORT

In **ational Application No**
PCT/CA 99/00449

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/645 A61K31/55

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 01157 A (THE WWK TRUST) 15 January 1998 (1998-01-15) page 1, line 4 - line 11 page 2, line 1 - page 3, line 7 page 4, line 11 - line 22 page 8, line 7 - line 19 page 11, line 23 - line 27 claims 1-5,8,9,12,13,16,19,20 --- -/--</p>	1-53

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

13 August 1999

Date of mailing of the international search report

01/09/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Economou, D

INTERNATIONAL SEARCH REPORT

In ternational Application No

PCT/CA 99/00449

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 97 10815 A (FROME, B.) 27 March 1997 (1997-03-27) claims 1,3,5,7,9,11,13,15,17,20,22 page 20, line 1 - page 31, paragraph 3 page 18, paragraph 3 - page 19, paragraph 1 page 17, last paragraph - page 18, line 3 page 16, last paragraph - page 17, paragraph 1 page 15, paragraphs 1,3 page 13, paragraph 3 page 11, last paragraph page 9 page 5, paragraph 4 page 1, line 1 - page 4, line 6</p>	1-53
X	<p>ORTEGA-ALVARO A. ET AL.: "The effects of different monoaminergic antidepressants on the analgesia induced by spinal cord adrenal medullary transplants in the formalin test in rats" ANESTHESIA AND ANALGESIA, vol. 84, no. 4, April 1997 (1997-04), pages 816-820, XP002112201 usa page 817, left-hand column - page 820, left-hand column, line 23 page 816, right-hand column abstract</p>	1-53
X	<p>SIERRALTA, F. ET AL.: "Effect of p-Chlorophenylalanine and alpha-Methyltyrosine on the Antinociceptive Effect of Antidepressant Drugs" PHARMACOLOGY AND TOXICOLOGY, vol. 77, no. 4, October 1995 (1995-10), pages 276-280, XP002112202 denmark page 277, right-hand column, line 10 - page 279, right-hand column, paragraph 5 page 277, left-hand column, paragraphs 4,6 page 276, left-hand column, paragraph 3 - page 276, right-hand column, paragraph 3 abstract</p>	1-53

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 99/00449

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9801157 A	15-01-1998	AU 3451797 A	02-02-1998
WO 9710815 A	27-03-1997	NONE	

THIS PAGE BLANK (USPTO)